



<b>Magellan Healthcare</b>	
<b>Clinical guidelines</b> <b>EXPERIMENTAL, UNPROVEN, OR</b> <b>INVESTIGATIONAL SERVICES</b>	<b>Original Date:</b> November 2, 2015 <b>Page 1 of 13</b>
<b>Physical Medicine – Clinical Decision Making</b>	<b>Last Review Date:</b> June 2017
<b>Guideline Number: NIA_CG_601</b>	<b>Last Revised Date:</b> June 2017
<b>Responsible Department:</b> <b>Clinical Operations</b>	<b>Implementation Date:</b> January 2018

### **Policy Statement**

This policy will be used to provide a listing of procedures considered experimental, investigational by any physical medicine practitioner. Services listed in the policy are not eligible for reimbursement.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

### **Purpose**

To provide a listing of procedures considered experimental, investigational or unproven services by any physical medicine practitioner, including chiropractors, physical therapists, occupational therapists, and speech language pathologists.

### **Scope**

Clinical Management, Coding, Customer Service, Claims and Contracting.

### **Coverage**

Coverage is subject to the terms of an enrollee’s benefit plan. To the extent there is any inconsistency between this medical policy and the terms of an enrollee’s benefit plan, the terms of the enrollee’s benefit plan documents will always control. Investigational services are not covered under enrollee’s health plan.

### **Definition**

A service is considered experimental/investigation if **any** of the following criteria is met:

1. The services, procedures or supplies requiring Federal or other Governmental body approval, such as drugs and devices, do not have unrestricted market approval from the Food and Drug Administration (FDA) or final approval from any other governmental regulatory body for use in treatment of a specified condition. Any approval that is granted as an interim step in the regulatory process is not a substitute for final or unrestricted market approval.
2. There is insufficient or inconclusive medical and scientific evidence to evaluate the therapeutic value of the service, procedure or supply.

3. There is inconclusive medical and scientific evidence in peer-reviewed medical literature that the service, procedure or supply has a beneficial effect on health outcomes.
4. The service, procedure or supply under consideration is not as beneficial as any established alternatives.
5. There is insufficient information or inconclusive scientific evidence that, when used in a non-investigational setting, the service, procedure or supply has a beneficial effect on health outcomes or is as beneficial as any established alternatives.

Experimental and investigational services include the use of a service, procedure or supply that is not recognized as standard clinical care for the condition, disease, illness or injury being treated. A service, procedure or supply includes, but is not limited to the diagnostic service, treatment, facility, equipment, or device. Magellan Healthcare and their client health plan will determine whether a service, procedure, or supply is considered experimental and investigational.

The following is a partial listing of experimental and investigational services:

- Advanced BioStructural Correction (ABC)
- Alphabiotics
- Applied Kinesiology or any of its derivations
- Applied Spinal Biomechanical Engineering
- BioEnergetic Synchronization Technique (B.E.S.T)
- Chiropractic Biophysics (CBP, Clinical Biomechanics of Posture, CBP Mirror Image Technique)
- Coccygeal Meningeal Stress Fixation
- Cold Laser Therapy
- Computerized muscle testing or analysis
- Craniosacral Therapy (CST)
- Directional Non-force Technique
- Spinal Diagnostic Ultrasound
- Hako-Med electrotherapy (horizontal electrotherapy)
- Hippotherapy
- Impulse adjusting instrument
- Intersegmental traction and Autotraction
- Kinesio taping (Elastic Therapeutic Taping)
- Live Cell Analysis or hair analysis
- Manipulation under Anesthesia (MUA)
- Moire Contourographic Analysis
- Nambudripad's Allergy Elimination Technique (NAET)/ other Allergy Testing
- National Upper Cervical Chiropractic Association (NUCCA technique)/Grostic technique
- Network Chiropractic, NeuroEmotional Technique (NET)
- Neurocalometer, Nervoscope, Nerve Conduction Velocity, Surface EMG, Paraspinal Electromyography, Spinoscopy or other nerve conduction testing for non-specific neck and back pain

- Neural Organizational Technique, Contact Reflex Analysis (CRA), Whole System Scan
- Nimmo Receptor-Tonus method
- Pettibon and wobble chair/board treatment
- Preventive Care, Maintenance Care, Corrective Care
- Pro-Adjuster
- Sacro Occipital Technique, Neurocranial Restructuring (NCR), Cranial Manipulation
- Sound Assisted Soft Tissue mobilization
- Chiropractic services directed at controlling progression and/or reducing scoliosis, including but not limited to the SpineCor brace and CLEAR scoliosis treatment
- Repeat imaging to determine the progress of conservative treatment
- Thermography
- Upledger Technique
- Vascular Studies, including, but not limited to, Doppler ultrasound analysis and plethysmography
- VAX-D, Lordex, LTX3000, DRX-9000, DRS (Decompression Reduction Stabilization System), or other back traction devices charged at a higher rate than mechanical traction (97012)
- Whole Body Vibration (WBV), Vibration Plate, Vibration Therapy
- Any lab work for which the office is not CLIA Certified or falls outside of the scope of practice, including, but not limited to: drug testing, therapeutic drug assays, and organ or disease oriented panels
- Treatment for brachioradial pruritis
- Dry Needling

Professional societies have published position statements concluding that diagnostic spinal ultrasound is investigational for non-operative spinal and paraspinal conditions in adults.

There is insufficient peer-reviewed published scientific evidence that computerized muscle testing leads to better patient outcomes. There is insufficient evidence to support any specific therapeutic effect of craniosacral therapy. While there is emerging evidence for the effectiveness of whole body vibration in treating some medical conditions, the evidence for whole body vibration as a treatment for LBP remains equivocal.

A 2015 systematic review found that that low level laser therapy is an effective method for relieving pain in non-specific chronic low back pain patients. However, no significant treatment effect was identified for disability scores or spinal range of motion outcomes. Yelden and colleagues (2009), concluded that there is no fundamental difference between LLLT and placebo LLLT when they are supplementing an exercise program for rehabilitation of patients with shoulder impingement syndrome. Ay and colleagues (2010), found no differences between laser and placebo laser treatments on pain severity and functional capacity in patients with acute and chronic low back pain caused by lumbar disc herniation. The Blue Cross and Blue Shield Association Technology Evaluation Center (2010) concluded that LLLT for either carpal tunnel syndrome or for chronic neck pain does not meet the Blue Cross and Blue Shield Association Technology Evaluation Center (TEC) criteria. Furthermore, the Work Loss Data Institute's clinical practice guideline on "Carpal tunnel syndrome" (2011) does not recommend LLLT as a therapeutic option. Kadhim-Saleh

et al (2013) examined the effectiveness of LLLT in reducing acute and chronic neck pain. The authors concluded that this systematic review provided inconclusive evidence because of significant between-study heterogeneity and potential risk of bias. They stated that the benefit seen in the use of LLLT, although statistically significant, does not constitute the threshold of minimally important clinical difference. Huang et al (2015), found that the best available current evidence does not support the effectiveness of low level laser therapy as a therapy for patients with knee osteoarthritis.

There is insufficient evidence to support the clinical value of the Pettibon System.

Thermography has not been shown to provide sufficient reliable characterizing information about neurologic dysfunction or deficit to accept it as a proven evaluative procedure for the clinical diagnosis or characterization of: neck or back pain; musculoskeletal pain; entrapment neuropathy; headache; or transient cerebral ischemia and stroke.

Magellan considers high-density surface electromyography (HD-sEMG), surface scanning EMG, paraspinal surface EMG, or macro EMG experimental and investigational as a diagnostic test for evaluating low back pain or other thoracolumbar segmental abnormalities such as soft tissue injury, intervertebral disc disease, nerve root irritation and scoliosis, and for all other indications because the reliability and validity of these tests have not been established. Surface EMG devices are also experimental and investigational for diagnosis and/or monitoring of nocturnal bruxism and all other indications because the reliability and validity of these tests have not been demonstrated. The Neurophysiologic Pain Profile (NPP) and the spine matrix scan (lumbar matrix scan) are considered experimental and investigational because the reliability and validity of these tests has not been established.

There is insufficient evidence to conclude that nerve conduction studies are beneficial for health outcomes in patients with non-specific neck or back pain. Non-invasive automatic or portable nerve conduction monitoring systems that test only distal motor latencies and conduction velocities are unproven and not medically necessary for the purpose of electrodiagnostic testing.

Plethysmography is used to diagnose deep vein thrombosis and arterial occlusive disease. Plethysmography is used as the sole diagnostic modality for these conditions or as an initial evaluation to determine the need for venography or arteriography. Body Plethysmography evaluates total lung capacity and residual volume. Since treatment of cardiovascular and lung conditions falls outside of the scope of chiropractic, patients should be referred for testing if these conditions are suspected.

## **Procedure:**

### **1. Guidelines:**

- a. If such services are to be provided, the practitioner will inform the member, in writing, that such services will be the member's responsibility. None of these services are to be performed in lieu of an appropriate examination or without consideration of an appropriate referral.

- b. There is limited scientific evidence that the use of experimental, investigational and unproven services provides an improved or more accurate diagnosis, nor do they result in an improved clinical outcome.
- c. Scientific literature will continue to be reviewed and any significant changes in published literature will be taken into consideration for modification of this policy.

**2. Exclusions/Limitations (not limited to):**

Refer to enrollee's Certificate of Coverage or Summary Plan Description.

**3. Removal of a service from the Experimental and Investigations Policy**

At least annually, a review of the current literature will be evaluated to determine if there is additional research in support of any of the services listed under this policy.

This evaluation will include the following criteria:

- Safety – Is the potential benefit superior to the potential harm?
- Health Outcomes – Is there evidence the service will provide, at minimum, equal outcomes and, at best, superior outcomes to currently available services?
- Patient Management - Will the service improve clinical decision making?
- Clinical Performance – Is the reliability as well as predictive value of the service equal or superior to the current “gold standard” for such services?
- Cost-effectiveness – Is the service equal to or lower cost than currently utilized services for similar diagnosis and treatment?

All criteria will be based on peer-reviewed scientific literature and internationally and nationally accepted and published guidelines. Peer-reviewed scientific studies must be published in or accepted for publication by medical journals meeting national requirements for scientific publication (<http://www.icmje.org>). The medical literature must meet the National Institutes of Health Library of Medicine for indexing (<http://www.nlm.nih.gov>). Medical journals that publish most of their scientific manuscripts by the editorial staff of a journal will not be considered for review. If the majority of funding for research is published by the device manufacturer or organization sponsoring a technique the results will not be considered for review.

If the service appears to be safe and cost-effective Magellan Healthcare will present these results to our health plan partners for consideration of coverage and/or payment. Final authority for such coverage determinations rests with the health plan.

## REFERENCES

- Aetna. (2013, July). Cold laser therapy [Clinical Policy Bulletin Number 363]. Hartford, CT: Aetna Inc.
- Aetna. (2014, March). Lumbar Traction Devices [Clinical Policy Bulletin Number 569]. Hartford, CT: Aetna Inc.
- Aetna. (2014, January). Manipulation Under Anesthesia [Clinical Policy Bulletin Number 204]. Hartford, CT: Aetna Inc.
- Aetna. (2014, February). Thermography [Clinical Policy Bulletin Number 29]. Hartford, CT: Aetna Inc.
- Aetna. (2013, August). Vertebral Axial Decompression Therapy [Clinical Policy Bulletin Number 180]. Hartford, CT: Aetna Inc.
- Aetna Clinical Policy Bulletin: Spinal Ultrasound – No. 0628. Accessed: 8.15.2016: [http://www.aetna.com/cpb/medical/data/600\\_699/0628.html](http://www.aetna.com/cpb/medical/data/600_699/0628.html)
- Albeck MJ, Taher G, et al. Diagnostic value of electrophysiological tests in patients with sciatica. *Acta Neurol Scand.* 2000;101(4):249-54.
- American Academy of Neurology. Review of the literature on spinal ultrasound for the evaluation of back pain and radicular disorders. *Neurology* 1998; 51:343–344.
- American Chiropractic College of Radiology. Policy statement. October, 1995. Adopted by ACA House of Delegates, December 1995.
- American College of Radiology. Ultrasound: Not effective in diagnosing spinal injuries. *ACR Bulletin.* Feb, 1996.
- American Institute of Ultrasound in Medicine. Official Statement: Diagnostic spinal ultrasound. Oct, 1995.
- Anrig CA. Spinal examination and specific spinal and pelvic adjustments. In Anrig CA, Plaugher G, eds. *Pediatric Chiropractic.* Williams and Wilkins, Baltimore. 1998:323-423.
- Aspegren DD, Wright RE, Hemler DE. Manipulation under epidural anesthesia with corticosteroid injection: two case reports. *J Manip Physiol Ther.* 1997;20(9):618-621.
- Barker WF, Gambale AG, Jackson BL. Reliability of the upper cervical x-ray marking system: A replication study. *Chiropr* 1988;1:10-13.
- Barrett S. Commercial hair analysis: science or scam? *JAMA.* 1974;229:1908-1909.
- Barry MS, et al. Facts and fallacies of diagnostic ultrasound of the adult spine. *Dynamic Chiropr.* 1996;14(9):30.

- Basford JR, Sheffield CG, Harmsen WS. Laser therapy: a randomized, controlled trial of the effects of low level intensity Nd:YAG laser irradiation on musculoskeletal back pain. *Arch Phys Med Rehabil.* 1999;80(6):647-52.
- Ben-David R, Raboy M. Manipulation under anesthesia combined with epidural steroid injection. *J Manip Physiol Ther.* 1994;17(9):605-609.
- Bendix T, Biering-Sorensen F. Posture of the trunk when sitting on forward inclining seats. *Scand J Rehabil Med.* 1983;15:1531-1535.
- Bertoti, DB. Effect of therapeutic horseback riding on posture in children with cerebral palsy. *Phys Ther.* 1988;68 1505-1512.
- Black KM, et al. The influence of different sitting positions on cervical and lumbar posture. *Spine.* 1996;21:65-70.
- Blomberg S, Tibblin G. A controlled, multicentre trial of manual therapy with steroid injections in low-back pain: Functional variables, side effects and complications during four months follow-up. *Clin Rehabil.* 1993;7(1):49-62.
- Blomberg S, Svardsudd K, Tibblin G. A randomized study of manual therapy with steroid injections in low-back pain. Telephone interview follow-up of pain, disability, recovery and drug consumption. *Eur Spine J.* 1994;3(5):246-254
- Bryner P, Cowin R. Hearing loss, otalgia and neck pain: A case report on long-term chiropractic care that helped to improve quality of life. *Chiropr J Aust* 2002;32:119-130.
- Cigna Medical Coverage Policy: Spinal Ultrasound – No. 0246. Accessed: 8.15.2016: [https://cignaforhcp.cigna.com/public/content/pdf/coveragePolicies/medical/mm\\_0246\\_coveragepositioncriteria\\_spine\\_ultrasound.pdf](https://cignaforhcp.cigna.com/public/content/pdf/coveragePolicies/medical/mm_0246_coveragepositioncriteria_spine_ultrasound.pdf)
- Clarke, J. A., et al., “Traction for Low-Back Pain With or Without Sciatica,” *Cochrane Database Syst Rev.*, Volume 2, 2007, CD003010.
- Colloca CJ, Keller TS. Electromyographic reflex responses to mechanical force manually assisted spinal manipulative therapy. *Spine.* 2001;26(10):1117-24.
- Cotchett MP, Munteanu SE, Landorf KB. Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Phys Ther* (published online prior to print) <http://ptjournalapta.org/content/early/2014/06/11/ptj.20130255>
- Davis GE, et al. A complication from neurocranial restructuring: nasal septum fracture. *Arch Otolaryn Head Neck Surg.* 2003;129:472-474.
- Debusse D, Chandler C, Gibb C. An exploration of German and British physiotherapists' views on the effects of hippotherapy and their measurement. *Physiother Theory Pract.* 2005;21(4):219-242.

- de Koning CH, van den Heuvel SP, Staal JB, et al. Clinimetric evaluation of methods to measure muscle functioning in patients with non-specific neck pain: A systematic review. *BMC Musculoskelet Disord*. 2008;9:142.
- Edzard Ernst MD, PhD, FMedSci, FSB, FRCP, FRCPEd. Craniosacral therapy: a systematic review of the clinical evidence. *Focus on Alternative and Complementary Therapies*, 17: 197–201.
- Federal Register, Centers for Medicare and Medicaid Services. Sensory Nerve Conduction Threshold Testing. Vol. 68, No. 143. July, 2003.
- Gammon SR, Mehlman CT, Chan W, Heifetz J, Durrett G, Wall EJ. A comparison of thoracolumbosacral orthoses and SpineCor treatment of adolescent idiopathic scoliosis patients using the Scoliosis Research Society standardized criteria. *J Pediatr Orthop*. 2010 Sep;30(6):531-8.
- Gutman, Gabriel et al. The effectiveness of the SpineCor brace for the conservative treatment of adolescent idiopathic scoliosis. Comparison with the Boston brace. *j.spinee*.2016.01.020, *The Spine Journal*, Volume 16, Issue 5, Pages 626–631.
- Furlan AD, van Tulder M, Cherkin D, et al. Acupuncture and dry-needling for low back pain: an updated systematic review with the framework of the Cochrane Collaboration. *Spine*. 2005; 30:944-963
- Gose EE, Naguszewski WK. Vertebral axial decompression therapy for pain associated with herniated or degenerated discs or facet syndrome. *J Neuro Res*. 1998;20:186-190.
- Green C, Martin CW, Bassett K, Kazanjian A. A systematic review of craniosacral therapy: biological plausibility, assessment reliability and clinical effectiveness. *Complement Ther Med*. 1999;7(4):201-7.
- Greenman PE. Manipulation with patient under anesthesia. *J Am Osteopath Assoc*. 1992;92:1159-70.
- Gruther W, Wick F, Paul B, et al. Diagnostic accuracy and reliability of muscle strength and endurance measurements in patients with chronic low back pain. *J Rehabil Med*. 2009;41(8):613-619.
- Gur A, Karakoc M, et al. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg Med*. 2003;32(3):233-8.
- Haldeman S, et al. Guidelines for Chiropractic Quality Assurance and Practice Parameters. Gaithersburg, MD:Aspen Publishers, Inc. 1993.
- Harrison DE, et al. The sacroiliac joint. *J Manipulative Physiol Ther*. 1997;20:607-617.

- Hartman SE, Norton JM. Interexaminer reliability and cranial osteopathy. *Scientific Review of Alternative Medicine*. 2002;6(1):23-34.
- Hawk C, Rupert R, Colonvega M, Boyd J, Hall S. Comparison of bioenergetic synchronization technique and customary chiropractic care for older adults with chronic musculoskeletal pain. *J Manip Physiol Ther* 2005;29(7):540-549.
- Hazell TJ, Olver TD, Hamilton CD, Lemon PW. Addition of synchronous whole-body vibration to body mass resistive exercise causes little or no effects on muscle damage and/or inflammation PW *J Strength Cond Res*. 2013 Apr 23
- Herzog J. Use of cervical spine manipulation under anesthesia for management of cervical disk herniation, cervical radiculopathy, and associated cervicogenic headache syndrome. *J Manip Physiol Ther*. 1999;22(3):166-170.
- Hochman JJ. Pierce and Pettibon combine their systems. *Today's Chiropr* 1991;20:84.
- Ireland TV, Pettibon BR, Morningstar MW, Schlappi H, Schilappi I. Reflex control of the spine and posture: a review of the literature from a chiropractic perspective. *Chiropr & Osteopat* 2005;13:34
- Jäkel A, von Hauenschild P. A systematic review to evaluate the clinical benefits of craniosacral therapy. *Complement Ther Med*. 2012 Dec;20(6):456-65
- Kenny JJ, Clemens R, et al. Applied kinesiology unreliable for assessing nutrient status. *J Am Dietetic Assoc*. 1988;88:698-704.
- Kietrys DM, Kerstin KM, Azzaretto E, et al. Effectiveness of dry needling for upper-quarter myofascial pain: a systematic review and meta-analysis. *J Orthop Sports Phys Ther*. 2013;43:620-634
- Kivmaki J, Pohjolainen T, Malmaivaara A, et al. Manipulation under anesthesia with home exercises versus home exercises alone in the treatment of frozen shoulder. A randomized controlled trial with 125 patients. *J Shoulder Elbow Surg*. 2007;16(6):722-726.
- Klevay LM, et al. Hair analysis in clinical and experimental medicine. *Am J Clin Nutrition*. 1987;46:233-236.
- Kohlbeck FJ, Haldeman S. Medication assisted spinal manipulation. *Spine*. 2002;2(4):288-302.
- Klougart N, Leboeuf-Yde C, Rasmussen LR. Safety in chiropractic practice. Part II: Treatment to the upper neck and the rate of cerebrovascular incidents. *J Manipulative Physiol Ther*. 1996;19(9):563-9.
- Labarre-Vila A. Surface electromyography procedure. *Rev Med Liege*. 2004;59:176-83.

- Lam FM, Lau RW, Chung RC, Pang MY. The effect of whole body vibration on balance, mobility and falls in older adults: a systematic review and meta-analysis. *Maturitas* 2012 Jul;72(3):206-13. doi: 10.1016/j.maturitas.2012.04.009. Epub 2012 May 18. The effect of WBV on other balance/mobility outcomes and fall rate remains inconclusive.
- Lindberg J, Carlsson J. The effects of whole-body vibration training on gait and walking ability - a systematic review comparing two quality indexes. There is low-quality evidence for WBV training having effects on gait and walking ability. Further research is needed. The evidence did not alter between the quality indexes. *Physiother Theory Pract* 2012 Oct;28(7):485-98. doi: 10.3109/09593985.2011.641670. Epub 2012 Jan 3.
- Ljunggren AE, Weber H, Larsen S. Autotraction versus manual traction in patients with prolapsed lumbar intervertebral discs. *Scand J Rehabil Med*. 1984;16(3):117-124.
- Lowell JA. Live cell analysis. *Nutrition Forum*. 1986;3:81-85.
- Merx JL, Thijssen HO, et al. Accuracy of ultrasonic evaluation of lumbar intervertebral discs by an anterior approach. *Neuroradiology*. 1989;31:386-90.
- Meyer JJ. The validity of thoracolumbar paraspinal scanning EMG as a diagnostic test: an examination of the current literature. *J Manipulative Physiol Ther*. 1994;17(8):539-51.
- Mohseni Bandpei, Rahmani N, Majdoleslam B, Abdollahi, Ahmad A. Reliability of surface electromyography in the assessment of paraspinal muscle fatigue: an updated systematic review. *J Manipulative Physiol Ther*. 2014 Sep;37(7):510-21.
- Monteforte P, Baratto L, et al. Low-power laser in osteoarthritis of the cervical spine. *Int J Tissue React*. 2003;25(4):131-6.
- Morningstar MW, Strauchman MN. Adolescent idiopathic scoliosis treatment using the Pettibon corrective procedures: a case report[letter]. *J Chiropr Med* 2007;6:83-84.
- Morningstar MW. Integrative treatment using chiropractic and conventional techniques for adolescent idiopathic scoliosis: Outcomes in four patients. *J Vert Sublux Res* 2007;9:1-7.
- Morningstar MW, Joy T. Scoliosis treatment using spinal manipulation and the Pettibon weighting system tm: a summary of 3 atypical presentations [case report]. *Chiropr & Osteopat* 2006;14:1-18.
- Morningstar MW. Improvement of lower extremity electrodiagnostic findings following a trial of spinal manipulation and motion-based therapy. 2006;14:6.
- Morningstar MW, Strauchman MN, Gilmour G. Adolescent idiopathic scoliosis treatment using Pettibon corrective procedures: A case report. *J Chiropr Med*. 2004;3:96-103.
- Morningstar MW. Cervical curve restoration and forward head posture reduction for the treatment of mechanical thoracic pain using the Pettibon corrective and rehabilitative procedures. *J Chiropr Med*. 2002;1:113-115.

- Nambudripad DS. Say goodbye to illness. Buena Park, CA. Delta Publishing Co, 1993.
- Negrini S, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Grivas TB, Kotwicki T, Maruyama T, Romano M, Vasiliadis ES. Braces for idiopathic scoliosis in adolescents. *Spine*. 2010;35(13):1285-93.
- Diagnostic ultrasound in sports medicine. Current concepts and advances. *Sports Med Arthrosc Rev*. 2009;17:25-30.
- Parreira PdCS, Costa LdCM, Hespanhol Junior LC, Lopes AD, Costa LOP. Current evidence does not support the use of Kinesio Taping in clinical practice: a systematic review. *Journal of Physiotherapy* 60: 31–39.
- Perraton L, Machotka Z, Kumar S. Whole-body vibration to treat low back pain: fact or fad? *Physiotherapy Canada*. *Physiotherapie Canada* [2011, 63(1):88-93]
- Pettibon BR. An introduction to spinal biomechanics. *Today's Chiropr* 1993;22:22-26.
- Pettibon B. Educating the insurance companies. *Today's Chiropr* 1989;18:74-75.
- Plaughner G. Skin temperature assessment for neuromusculoskeletal abnormalities of the spinal column. *J Manipulative Physiol Ther*. 1992;15(6):365-81.
- Pullman SL, et al. Clinical utility of surface EMG: report of the therapeutics and technology assessment subcommittee of the American Academy of Neurology. *Neurology*. 2000;55:171-77.
- Ramos G, Martin W. Effects of vertebral axial decompression on intradiscal pressure. *J Neurosurg*. 1994;81:35-353.
- Seidel S, et al. Assessment of commercial laboratories performing hair mineral analysis. *JAMA*. 2001;285:67-72.
- Schuster TL, Dobson M, et al. Wellness lifestyles II: modeling the dynamic of wellness, health lifestyle practices and Network Spinal Analysis. *J Altern Complement Med*. 2004;10(2):357-67.
- Seferlis T, Nemeth G, Carlsson AM, Gillstrom P. Conservative treatment in patients sick-listed for acute low-back pain: A prospective randomised study with 12 months' follow-up. *Eur Spine J*. 1998;7(6):461-470.
- Silkwood-Scherer D, Killian C, Long TM, Martin KS. Hippotherapy-an intervention to habilitate balance deficits in children with movement disorders. *Physical Therapy* 2012; 92(5):707-717.
- Sitja Rabert M, Rigau Comas D, Fort Vanmeerhaeghe A, Santoyo Medina C, Rogue FM, Romero-Rodríguez D, Bonfill Cosp X. Whole-body vibration training for patients

with neurodegenerative disease. There is insufficient evidence of the effect of WBV training on functional performance of neurodegenerative disease patients. Also, there is insufficient evidence regarding its beneficial effects on signs and symptoms of the disease, body balance, gait, muscle strength and quality of life compared to other active physical therapy or passive interventions in Parkinson's disease or multiple sclerosis. *Cochrane Database Syst Rev.* 2012 Feb 15;2:CD009097. doi: 10.1002/14651858.CD009097.pub2.

Snider L, Korner-Bitensky N, Kammann C, et al. Horseback riding as therapy for children with cerebral palsy: Is there evidence of its effectiveness? *Phys Occup Ther Pediatr.* 2007;27(2):5-23.

Spector B, et al. Manual of procedures for Moire contourography. New York Chiropractic College, 1979.

Staeble HJ, Koch MJ, Pioch T. Double-blind study on materials testing with applied kinesiology. *J of Dental Research.* 2005;84(11):1066-1069.

Sterba JA. Does horseback riding therapy or therapist-directed hippotherapy rehabilitate children with cerebral palsy? *Dev Med Child Neurol.* 2007;49(1):68-73.

Tilaro F. An overview of vertebral axial compression. *Canad J Clin Med.* 1998;5:1-7.

Tesio L, Merlo A. Autotrraction versus passive traction: An open controlled study in lumbar disc herniation. *Arch Phys Med Rehabil.* 1993;74(8):871-876.

The American Institute for Ultrasound in Medicine. Official Statement: Nonoperative spinal/paraspinal ultrasound in adults. April 6, 2009. Accessed: 8.15.2016  
<http://www.aium.org/officialStatements/18>

Troyanovich SJ. Motion palpation it's time to accept the evidence. *J Manipulative Physiol Ther.* 1998;21:568-571.

Tullberg T, Svanborg E, et al. A preoperative and postoperative study of the accuracy and value of electrodiagnosis in patients with lumbosacral disc herniation. *Spine.* 1993;18(7):837-42.

Tough EA, White AR. Effectiveness of acupuncture/dry needling for myofascial trigger point pain. *Phys Ther Rev.* 2011;2: 147-54

UnitedHealthcare Medical Policy: Spinal Ultrasonography – No. 2010T0462E. Accessed 8.15.2016: [www.unitedhealthcareonline.com/](http://www.unitedhealthcareonline.com/)

van den Noort S, et al. Assessment: Thermography in Neurologic Practice. *Neurology* 1990; 40:523-5

VAX-D: No evidence from controlled trials to support claims of efficacy. *The Back Letter.* 1998;13(9):97, 104, 105.

Versendaal DA. Contact reflex analysis and designed clinical nutrition. Jenison, MI:Hoezee Marketing, 1993.

Weiss HR. SpineCor vs. natural history - explanation of the results obtained using a simple biomechanical model. Stud Health Technol Inform. 2008;140:133-6.

Wong MS, Cheng JC, Lam TP, Ng BK, Sin SW, Lee-Shum SL, Chow DH, Tam SY. The effect of rigid versus flexible spinal orthosis on the clinical efficacy and acceptance of the patients with adolescent idiopathic scoliosis. Spine. 2008;33(12):1360-5.

Williams S, Whatman C, Hume PA, Sheerin K. Kinesio taping in treatment and prevention of sports injuries: a meta-analysis of the evidence for its effectiveness. Sports Med 2012;42(2):153-64.

Zazarian LN, et al. Paraspinal ultrasonography: Lack of accuracy in evaluating patients with cervical or lumbar back pain. J Ultrasound Med. 1998;17:117-22.

Reviewed/Approved by *Michael Pentecost MD* Michael Pentecost, MD, Chief Medical Officer