

Heart Catheterization

Clinical Guideline Tip Sheet

*This tip sheet is intended to further assist you with the clarification of the Magellan Healthcare¹ clinical guidelines. It is for informational purposes only and is **NOT** intended as a substitute for the clinical guidelines that must be utilized when reviewing cases for medical necessity and clinical appropriateness.*

Overview

Documentation/reports/testing needs to be provided for review of request. Office notes provided should explain the plan for arteriography, based on an increasing pattern of typical symptoms of a concern for Unstable angina.

Recommendations

Stable Ischemic Heart Disease:

- ❖ Symptoms of ischemia and cannot undergo stress testing or CCTA, and there is a high likelihood the outcome will affect therapy
- ❖ ETT with high risk DTS (-11) or ST elevation, hypotension or VT during exercise or several minutes of ST depression persisting into recovery
- ❖ Low risk stress imaging with new or worsening symptoms concerning for coronary origin despite optimal medical therapy (OMT) or documentation that patient cannot tolerate OMT
- ❖ Intermediate risk findings on stress imaging - (see guideline background section) with symptoms suggestive of CAD, unsatisfactory quality of life due to angina symptoms or EF less than 50%
- ❖ High risk findings on stress imaging including:
 - Resting left ventricular (LV) dysfunction (LVEF <35%)
 - Severe stress-induced left ventricular (LV) dysfunction
 - Stress-induced perfusion abnormalities ≥10% myocardium or stress indicating multiple vascular territories with abnormalities
 - Stress-induced LV dilation
- ❖ Discordant/Inconclusive non-invasive results in symptomatic patients (i.e.- strongly positive stress ECG portion with low risk imaging)

¹National Imaging Associates, Inc. is an affiliate of Magellan Healthcare, Inc.

CCTA Abnormalities:

- ❖ Symptomatic patients with hemodynamically significant disease (70% or more) on CCTA
- ❖ Symptomatic patients with obstructive disease (between 50-69%) in 2 or more vessels
- ❖ Left main stenosis of 50% or greater (regardless of symptoms)

Heart Failure and Left Ventricular Dysfunction (in patients who are candidates for coronary revascularization)

- ❖ Newly recognized Reduction in EF to $\leq 50\%$ with intermediate risk stress test findings (see guidelines for definition) and symptoms of ischemia
- ❖ New wall motion abnormality and symptoms suggestive of ischemia
- ❖ To investigate structural heart disease when there is a concern for ischemic etiology (secondary MR/VSD)
- ❖ To investigate etiology of diastolic heart failure where there is reasonable likelihood of CAD (based on symptoms or imaging studies)

Ventricular Arrhythmias (without identified non-cardiac cause)

- ❖ Recovery post cardiac arrest
- ❖ Sustained ventricular tachycardia or ventricular fibrillation
- ❖ Exercise induced Ventricular tachycardia

Prior to non-coronary cardiac surgery (i.e. prior to valve replacement, repair of aneurysm) in a patient with:

- ❖ Symptoms of angina
- ❖ History of CAD or with cardiac risk factors (includes men > 40 or post-menopausal women)
- ❖ LV function $\leq 50\%$
- ❖ Prior to TAVR
- ❖ Non-invasive data that shows objective evidence of ischemia
- ❖ When more detailed assessment of coronary artery anatomy (including anomalous origins) is necessary

Post Cardiac Transplantation:

- ❖ Assessment for annual graft vasculopathy for the first 5 years, followed by annual assessment if there is documented allograft vasculopathy
- ❖ Any clinical change (new LV dysfunction, ischemic symptoms, non-invasive findings of ischemia)

Hemodynamic Assessment Magellan Healthcare does not manage right heart catheterization as a stand-alone procedure)

- ❖ Discordance between non-invasive data and clinical picture when management will be changed by the results of the angiogram
- ❖ Hemodynamic assessment of bio prosthetic or mechanical valve when TTE and TEE images are inadequate and CMR or CCT are not readily available

References

A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *JACC*. 2012; 59(22):1995-2027.

Al-Khatib SM, Stevenson WG, Ackerman MUJ, et al. 2017 AHA/ACC/HRS Guideline for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death. Accepted Manuscript in Press. *JACC*. Available at:

https://www.jacc.org/doi/pdf/10.1016/j.jacc.2017.10.054?_ga=2.50014127.860733034.1652283446-1800485333.1648040047 Retrieved May 1, 2018

Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 Multimodality Appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: A report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *JACC*. 2014;63(4):380-406. Available at:

https://www.jacc.org/doi/pdf/10.1016/j.jacc.2013.11.009?_ga=2.212610333.860733034.1652283446-1800485333.1648040047

2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease

Karen K. Stout, Curt J. Daniels, Jamil A. Aboulhosn, Biykem Bozkurt, Craig S. Broberg, Jack M. Colman, Stephen R. Crumb, Joseph A. Dearani, Stephanie Fuller, Michelle Gurvitz, Paul Khairy, Michael J. Landzberg, Arwa Saidi, Anne Marie Valente, George F. Van Hare

Journal of the American College of Cardiology Aug 2018, 25255; DOI: 10.1016/j.jacc.2018.08.1029

Doherty JU, Kort S, Mehran R. et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2017 Appropriate Use Criteria for Multimodality Imaging in Valvular Heart Disease. A Report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *JACC*. 2017; 70(13):1647-1672.

