

National Imaging Associates, Inc. *	
Clinical guidelines: HODGKIN LYMPHOMA	Original Date: June 2013
Radiation Oncology	Last Revised Date: February 2021
Guideline Number: NIA_CG_132	Implementation Date: January 2022

INDICATIONS FOR RADIATION THERAPY AND TREATMENT OPTIONS

2D and 3D conformal radiation therapy techniques are considered medically necessary for treatment of Hodgkin's Lymphoma (NCCN, 2021)

Stage I-II (nonbulky disease)

- Chemotherapy + radiation therapy (20-30 Gy) up to 20 fractions

Stage IB-IIB (nonbulky disease)

- Chemotherapy + radiation therapy (30 Gy) up to 20 fractions

Stage I-IV (bulky disease)

- Chemotherapy + radiation therapy (30-36 Gy) up to 24 fractions

Palliative

- Up to 10 fractions of external radiation may be indicated for symptom control

Radiation therapy alone is uncommon (except for lymphocyte predominant Hodgkin lymphoma). If used, doses of 30-36 Gy (up to 20 fractions) is recommended for uninvolved regions, 25-30 Gy (up to 17 fractions)

TREATMENT OPTIONS REQUIRING PHYSICIAN REVIEW

Intensity Modulated Radiation Therapy (IMRT)

IMRT is not indicated as a standard treatment option and should not be used routinely for the delivery of radiation therapy for Hodgkin's lymphoma. IMRT is strictly defined by the utilization of inverse planning modulation techniques. IMRT may be appropriate for limited circumstances in which radiation therapy is indicated and 3D conformal radiation therapy (3D-CRT) techniques cannot adequately deliver the radiation prescription without exceeding normal tissue radiation tolerance, the

* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

delivery is anticipated to contribute to potential late toxicity or tumor volume dose heterogeneity is such that unacceptable hot or cold spots are created.

Clinical rationale and documentation for performing IMRT rather than 2D or 3D-CRT treatment planning and delivery will need to:

- Demonstrate how 3D-CRT isodose planning cannot produce a satisfactory treatment plan (as stated above) via the use of patient-specific dose volume histograms and isodose plans.
- Provide tissue constraints for both the target and affected critical structures.

Stereotactic Body Radiation Therapy

Stereotactic Body Radiation Therapy (SBRT) is not currently a routine treatment option for the treatment of Hodgkin's lymphoma. SBRT may be appropriate for patients with tumors arising in or near a previously irradiated region to minimize risk to surrounding normal tissues. (ASTRO,2014). If requested, this would require peer to peer review to determine medical necessity.

Proton Beam Radiation Therapy

Proton beam is not an approved treatment option for Hodgkin's Lymphoma. Proton beam has not been proven superior treatment to conventional radiation therapy.

THE FOLLOWING APPLIES TO CMS (MEDICARE) MEMBERS ONLY

For Proton Beam and Stereotactic Radiotherapy, refer to Local Coverage Determination (LCD), if applicable.

BACKGROUND

Due to the significant improvement in treatment for this disease, Hodgkin disease is further classified into classical Hodgkin lymphoma (that accounts for 95% of all Hodgkin cases) and lymphocyte predominant Hodgkin lymphoma. Staging for Hodgkin lymphoma is based on the Ann Arbor staging system (stage I-IV), further subdivided into "A" (no systemic symptoms presents) and "B" (weight loss of >10%, fevers, or night sweats). Unfavorable prognostic factors include bulky mediastinal disease, nodal mass >10 cm, numerous sites of disease, significantly elevated erythrocyte sedimentation rate, or B symptoms. Treatment recommendations are typically based on three subgroups of Hodgkin lymphoma: early stage favorable (stage I-II with no unfavorable factors), early stage unfavorable (stage I-II with any unfavorable factors as mentioned above), and advanced stage disease (stage III and IV). When radiation therapy is used for the treatment of Hodgkin disease, it is usually in combination with chemotherapy. If chemotherapy is used alone, radiation therapy can be used for relapse.

POLICY HISTORY

Date	Summary
February 2021	Added content to clarify SBRT
February 2020	Updated references
February 2019	Added and updated references

REFERENCES

- Abuzetun JY, Loberiza F, Vose J, et al. The Stanford V regimen is effective in patients with good risk Hodgkin lymphoma but radiotherapy is a necessary component. *Br J Haematol*. 2009; 144:531-537. doi: 10.1111/j.1365-2141.2008.07500.x.
- Advani RH, Hoppe RT, Baer DM, et al. Efficacy of abbreviated Stanford V chemotherapy and involved field radiotherapy in early stage Hodgkin's disease: Mature results of the G4 trial. *Blood*. 2009; 114:1670. doi: 10.1093/annonc/mds542.
- Aleman BM, Raemaekers JM, Tomisic R, et al. Involved-field radiotherapy for patients in partial remission after chemotherapy for advanced Hodgkin's lymphoma. *Int J Radiat Oncol Biol Phys*. 2007; 67:19-30. doi:10.1016/j.ijrobp.2006.08.041.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Hodgkin Lymphoma – Favorable Prognosis Stage I and II. <https://acsearch.acr.org/docs/69352/Narrative/>. Reviewed 2016. Accessed April 27, 2018.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Hodgkin Lymphoma – Stage III and IV. <https://acsearch.acr.org/docs/69348/Narrative/>. Reviewed 2016. Accessed April 27, 2018.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Hodgkin Lymphoma – Unfavorable Clinical Stage I and II. <https://acsearch.acr.org/docs/69389/Narrative/>. Reviewed 2015. Accessed May 11, 2016.
- American Society for Radiation Oncology (ASTRO). Model Policies. Stereotactic Body Radiation Therapy. ASTRO; 2014. Retrieved from: https://www.astro.org/uploadedFiles/_MAIN_SITE/Daily_Practice/Reimbursement/Model_Policies/Content_Pieces/ASTROSBRTModelPolicy.pdf. Accessed May 3, 2018.
- Aversa SM, Salvagno L, Sorarù M, et al. Stanford V regimen plus consolidative radiotherapy is an effective therapeutic program for bulky or advanced-stage Hodgkin's disease. *Acta Haematol*. 2004; 112:141-147. doi: 10.1159/000079725.
- Bentzen SM, Constine LS, Deasy JO, et al. Quantitative analyses of normal tissue effects in the clinic QUANTEC: An introduction to the scientific issues. Introductory paper. *Int J Radiat Oncol Biol Phys*. 2010; 76(3):S3-S9.
- Bonadonna G, Bonfante V, Viviani S, et al. ABVD plus subtotal nodal versus involved-field radiotherapy in early-stage Hodgkin's disease: Long-term results. *J Clin Oncol*. 2004; 22: 2835-2841. doi: 10.1200/JCO.2004.12.170.
- Canellos GP, Abramson JS, Fisher DC, et al. Treatment of favorable, limited-stage Hodgkin's lymphoma with chemotherapy without consolidation by radiation therapy. *J Clin Oncol*. 2010; 28:1611-1615. doi:

10.1200/JCO.2009.25.3260.

Cella L, Liuzzi R, Magliulo M, et al. Radiotherapy of large target volumes in Hodgkin's lymphoma: Normal tissue sparing capability of forward IMRT versus conventional techniques. *Radiat Oncol*. May 11, 2010; 5:33.

Chera BS, Rodriguez C, Morris CG, et al. Dosimetric comparison of three different involved nodal irradiation techniques for stage II Hodgkin's lymphoma patients: Conventional radiotherapy, intensity-modulated radiotherapy, and three-dimensional proton radiotherapy. [Published online ahead of print April 20, 2009]. *Int J Radiat Oncol Biol Phys*. November 15, 2009; 75(4):1173-80.

Edwards-Bennett SM, Jacks LM, Moskowitz CH, et al. Stanford V program for locally extensive and advanced Hodgkin lymphoma: The Memorial Sloan-Kettering Cancer Center experience. *Ann Oncol*. 2010; 21:574-581. doi: 10.1093/annonc/mdp337.

Eich HT, Diehl V, Gørgen H, et al. Intensified chemotherapy and dose-reduced involved-field radiotherapy in patients with early unfavorable Hodgkin's lymphoma: Final analysis of the German Hodgkin Study Group HD11 trial. *J Clin Oncol*. 2010; 28: 4199-4206. doi: 10.1200/JCO.2010.29.8018.

Engert A, Franklin J, Eich HT, et al. Two cycles of doxorubicin, bleomycin, vinblastine, and dacarbazine plus extended-field radiotherapy is superior to radiotherapy alone in early favorable Hodgkin's lymphoma: Final results of the GHSG HD7 trial. *J Clin Oncol*. 2007; 25:3495-3502. doi: 10.1200/JCO.2006.07.0482.

Engert A, Plütschow A, Eich HT, et al. Reduced treatment intensity in patients with early-stage Hodgkin's lymphoma. *N Engl J Med*. 2010; 363:640-652. doi: 10.1056/NEJMoa1000067.

Fermé C, Eghbali H, Meerwaldt JH, et al. Chemotherapy plus involved-field radiation in early-stage Hodgkin's disease. *N Engl J Med*. 2007; 357:1916-1927. doi: 10.1056/NEJMoa064601.

Gordon LI, Hong F, Fisher RI, et al. A randomized phase III trial of ABVD vs. Stanford V +/- radiation therapy in locally extensive and advanced-stage Hodgkin Lymphoma: An intergroup study coordinated by the Eastern Cooperative Oncology Group (E2496). [Published online ahead of print November 26, 2012]. *J Clin Oncol*. February 20, 2013; 31(6):684-691.

Hoppe RT. Hodgkin's lymphoma: The role of radiation in the modern combined strategies of treatment. *Hematol Oncology Clin North Am*. 2007; 21:915-927. <http://www.ncbi.nlm.nih.gov/pubmed/17908628>.

Hoskin PJ, Lowry L, Horwich A, et al. Randomized comparison of the stanford V regimen and ABVD in the treatment of advanced Hodgkin's lymphoma: United Kingdom National Cancer Research Institute Lymphoma Group Study ISRCTN 64141244. *J Clin Oncol*. 2009; 27:5390-5396. doi: 10.1200/JCO.2009.23.3239.

Jackson A, Marks LB, Bentzen SM, et al. The lessons of QUANTEC: Recommendations for reporting and gathering data on dose-volume dependencies of treatment outcome. *Int J Radiat Oncol Biol Phys*. 2010; 76(3):S155-S160.

Jacobson CA, Longo DL. Management of early-stage Hodgkin's lymphoma. *Principles and Practice of Oncology*. 2011; 25(2):2011.

Johnson PW, Sydes MR, Hancock BW, et al. Consolidation radiotherapy in patients with advanced Hodgkin's lymphoma: Survival data from the UKLG LY09 randomized controlled trial (ISRCTN97144519). *J Clin Oncol*. 2010; 10:3352-3359. doi: 10.1200/JCO.2009.26.0323.

Koeck J, Abo-Madyan Y, Lohr F, et al. Radiotherapy for early mediastinal Hodgkin lymphoma according to the German Hodgkin Study Group (GHSG): The roles of intensity-modulated radiotherapy and involved-node radiotherapy. *Int J Radiat Oncol Biol Phys*. May 1, 2012; 83(1):268-276.

Li J, Dabaja B, Reed V, et al. Rationale for and preliminary results of proton beam therapy for mediastinal lymphoma. *Int J Radiat Oncol Biol Phys*. 2011; 81:167-174. doi: 10.1016/j.ijrobp.2010.05.007.

Macdonald DA, Connors JM. New strategies for the treatment of early stages of Hodgkin's lymphoma. *Hematol Oncol Clin North Am*. 2007; 21:871-880. <http://dx.doi.org/10.1016/j.hoc.2007.06.014>.

Meyer RM, Gospodarowicz MK, Connors JM, et al. Randomized comparison of ABVD chemotherapy with a strategy that includes radiation therapy in patients with limited-stage Hodgkin's lymphoma: National Cancer Institute of Canada Clinical Trials Group and the Eastern Cooperative Oncology Group. *J Clin Oncol*. 2005; 23:4634-4642. doi: 10.1200/JCO.2005.09.085.

National Comprehensive Cancer Network (NCCN). Hodgkin Lymphoma V2.2021. [_https://www.nccn.org/professionals/physician_gls/pdf/hodgkins.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hodgkins.pdf). Accessed on Feb 11, 2021.

Paumier A, Ghalibafian M, Gilmore J, et al. Dosimetric benefits of intensity-modulated radiotherapy combined with the deep-inspiration breath-hold technique in patients with mediastinal Hodgkin's lymphoma. [Published online ahead of print June 24, 2011]. *Int J Radiat Oncol Biol Phys*. March 15, 2012; 82(4):1522-1527.

Paumier A, Ghalibafian M, Beaudre A, et al. Involved-node radiotherapy and modern radiation treatment techniques in patients with Hodgkin lymphoma. *Int J Radiat Oncol Biol Phys*. May 1, 2011; 80(1):199-205.

Rueda DA, Márquez A, Gumá J, et al. Treatment of stage I and II Hodgkin's lymphoma with ABVD chemotherapy: Results after 7 years of a prospective study. *Ann Oncol*. 2004; 15:1798-1804. doi: 10.1093/annonc/mdh465.

Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin*. January/February 2012;

62(1):10-29. doi: 10.3322/caac.20138.

Specht L, Yahalom J, Illidge T, et al. Modern radiation therapy for Hodgkin lymphoma: Field and dose guidelines from the international lymphoma radiation oncology group (ILROG). *Int J Radiat Oncol Biol Phys*. July 15, 2014. <http://www.ncbi.nlm.nih.gov/pubmed/23790512>. Accessed May 26, 2016.

Weber DC, Johanson S, Peguret N, et al. Predicted risk of radiation-induced cancers after involved field and involved node radiotherapy with or without intensity modulation for early-stage hodgkin lymphoma in female patients. [Published online ahead of print August 26, 2010]. *Int J Radiat Oncol Biol Phys*. October 1, 2011; 81(2):490-497.

Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.