

# ONCOLOGY: ALGORITHMIC TESTING

---

## OVERVIEW

Oncology prognostic and algorithmic tests are developed to aid in determining the likelihood that an individual has cancer, the prognosis for a patient diagnosed with cancer, and/or surveillance for recurrence. These tests may be used to guide clinical decision making for an individual diagnosed with cancer. The testing methodologies include Gene Expression Profiling (GEP), which analyzes messenger RNA (mRNA) typically of multiple genes simultaneously, multimarker serum analysis, single-nucleotide variant testing, plasma-based proteomic analysis, and incorporation of other clinical data into test outputs.

In addition to the tests previously mentioned, proteogenomic testing is an emerging area. Proteogenomic testing combines the analysis of DNA with RNA and/or protein analysis. The current focus of proteogenomics is primarily on diagnostic and prognostic analyses in various cancers. Results also seek to provide potential treatment options, and to which treatments the cancer may be resistant.

Polygenic Risk Score (PRS) tests are another emerging area. These tests combine information from population SNP analysis with clinical and family history and aim to give additional insight into an individual's lifetime risk to develop a specific cancer.

Results of prognostic and algorithmic tests are often reported as a recurrence score, probability of distant disease recurrence, malignant potential, probable site of origin, or cancer risk score. Additionally, the output of these prognostic and algorithmic tests may be useful to assist in surgical and management decision-making and to identify individuals who may benefit from adjuvant therapy.

## POLICY REFERENCE TABLE

Below is a list of higher volume tests and the associated laboratories for each coverage criteria section. This list is not all inclusive

<a href="#">Coverage Criteria Sections</a>	<b>Example Tests, Labs</b>	<b>Common CPT Codes</b>	<b>Common ICD Codes</b>	<a href="#">Ref</a>
<b>Breast Cancer</b>				
<a href="#">Breast Cancer Treatment and Prognostic Algorithmic Tests</a>	Oncotype Dx Breast Recurrence Score (Exact Sciences)	81519, S3854	C50.011-C50.92, Z17.0	1
	Breast Cancer Index Prognostic (bioTheranostics)	81518, S3854		
<a href="#">Breast Cancer Prognostic Algorithmic Tests</a>	EndoPredict (Myriad)	81522, S3854	C50, Z17.0, Z17.1	1
	MammaPrint (Agendia, Inc.)	81521, 81523 S3854		
	Prosigna Assay (NeoGenomics)	81520		
<a href="#">Gene Expression Profiling Breast Cancer Subtyping Tests</a>	BluePrint (Agendia, Inc.)	81599, S3854	C50-C50.929	1
	Insight TNBCtype™ (Insight Molecular Labs)	0153U		
<a href="#">Breast DCIS Prognostic Algorithmic Tests</a>	Oncotype DX Breast DCIS Score (Exact Sciences)	0045U	D05.1	1
<b>Colorectal Cancer</b>				
<a href="#">Colorectal Cancer Prognostic Algorithmic Tests</a>	Oncotype DX Colon Recurrence Score (Exact Sciences)	81525	C18.0-C18.9	2
	miR-31now (GoPath Laboratories)	0069U		
<b>Prostate Cancer</b>				
<a href="#">Prostate Cancer Treatment and Prognostic Algorithmic Tests</a>	Oncotype DX Genomic Prostate Score (MDxHealth)	0047U	C61	3, 19
	Decipher Prostate Biopsy Genomic Classified Classifier (Veracyte)	81542		
	Decipher Prostate RP			

	Genomic Classifier (Veracyte)			
	Prolaris (Myriad Genetics)	81541		
<a href="#">Prostate Cancer Risk Assessment Algorithmic Tests</a>	4K Prostate Score (Serum) (BioReference Laboratories)	81539	C61, Z12.5	4, 27, 29
	Prostate Health Index (ARUP Laboratories)	84153, 84154, 86316		
	SelectMDx for Prostate Cancer (MDx Health)	0339U		
	ExoDx Prostate Test (ExosomeDx)	0005U		
	IsoPSA® (Cleveland Diagnostics, Inc)	0359U		
<a href="#">Prostate Cancer Diagnostic Algorithmic Tests</a>	ConfirmMDx for Prostate Cancer (MDxHealth)	81551	C61, Z12.5	5, 29
	MyProstateScore (MPS) (University of Michigan MLabs)	0113U		
<b><a href="#">Thyroid Cancer</a></b>				
<a href="#">Thyroid Cancer Diagnostic Algorithmic Tests</a>	ThyroSeq Genomic Classifier (CBLPath)	0026U	C73, D44.0, E04.1	6, 7, 8
	ThyGeNEXT (Interpace Diagnostics)	0245U		
	ThyraMIR (Interpace Diagnostics)	0018U		
	Afirma Genomic Sequencing Classifier (Veracyte)	81546		
	Afirma Xpression Atlas (Veracyte)	0204U		
<b><a href="#">Uveal Melanoma</a></b>				
<a href="#">Uveal Melanoma Prognostic Algorithmic Tests</a>	DecisionDX-UM (Castle Bioscience, Inc.)	81552	C69	9
<b><a href="#">Cutaneous Melanoma</a></b>				

<a href="#">Cutaneous Melanoma Prognostic Algorithmic Tests</a>	DecisionDX-Melanoma (Castle Biosciences, Inc.)	81529	C43, D03.0-D03.9, Z12.83	10, 11
<a href="#">Cutaneous Melanoma Diagnostic Algorithmic Tests</a>	myPath Melanoma (Castle Biosciences Inc)	0090U	D22.0-D22.9, D48.5, D49.2, Z12.83	10, 11, 26
<a href="#">Cutaneous Melanoma Risk Assessment Algorithmic Tests</a>	Pigmented Lesion Assay (DermTech)	0089U	D22-D23, Z12.83	23
<b>Ovarian Cancer</b>				
<a href="#">Ovarian Cancer Diagnostic Algorithmic Tests</a>	OVA1 (Aspira)	81503	D27.0, D27.1, D27.9, D39.10-D39.12, D39.9, D49.59, D49.9	12
	Overa (Aspira)	0003U		
	Ovarian Malignancy Risk (ROMA) (LabCorp)	81500		
	OvaWatch (Aspira Women's Health)	0375U		
<a href="#">Ovarian Cancer Treatment Algorithmic Tests</a>	myChoice CDx (Myriad Genetics)	0172U	C48, C56, C57.0	12, 20
<b>Gynecologic Cancer</b>				
<a href="#">Gynecologic Cancer Treatment Algorithmic Tests</a>	ChemoFx (Helomics Corporation)	81535	C51-C57	12, 17, 18
	ChemoFx - Additional Drug (Helomics Corporation)	81536		
<b>Lung Cancer</b>				
<a href="#">Lung Cancer Diagnostic Algorithmic Tests</a>	Nodify XL2 (Biodesix)	0080U	R91.1	25
	REVEAL Lung Nodule Characterization (MagArray)	0092U		
	Percepta Bronchial Genomic Classifier (Veracyte)	81599		
<a href="#">Lung Cancer</a>	VeriStrat (Biodesix)	81538	C34, D38.1,	24

<a href="#">Treatment Algorithmic Tests</a>	DetermaRx (Oncocyte)	0288U	D38.6	
<b>Bladder and Urinary Tract Cancer</b>				
<a href="#">Bladder Cancer Diagnostic and Recurrence Algorithmic Tests</a>	Cxbladder Triage (Pacific Edge)	0363U	C67, D09.0, D49.4, R31.9, Z85.51	13, 14
	Cxbladder Detect (Pacific Edge)	0012M		
	Cxbladder Monitor (Pacific Edge)	0013M		
	Oncuria Detect (DiaCarta Clinical Lab)	0365U		
	Oncuria Monitor (DiaCarta Clinical Lab)	0366U		
	Oncuria Predict (DiaCarta Clinical Lab)	0367U		
<a href="#">Urinary Tract Cancer Recurrence Algorithmic Tests</a>	Alere NMP22® (Alere)	86386	C67 D09.0, D49.4, R31.9, Z85.51	
	Alere NMP22® BladderChek® (Alere)	86386		
<b>Pancreatic Cancer</b>				
<a href="#">Pancreatic Cyst Risk Assessment Algorithmic Tests</a>	PancaGEN (Interpace Diagnostics)	81479	D49, K86.2	21, 22
	Pancreatic Cyst Fluid NGS Analysis-PancreaSeq (Univ of Pittsburgh Medical Center)	0313U		
<b>Cancer of Unknown Primary</b>				
<a href="#">Cancer of Unknown Primary Gene Expression Profiling Tests</a>	CancerTYPE ID (Biotheranostics)	81540	C79.9, C80.0, C80.1	16
<b>Polygenic Risk Score Tests</b>				
<a href="#">Breast Cancer</a>	BrevaGenplus (Pathogen)	81599	Z13.71, Z13.79	15

<a href="#">Polygenic Risk Score Tests</a>	Sciences Laboratories)		Z80.3	
<a href="#">Multiple Myeloma Polygenic Risk Score Tests</a>	Myeloma Prognostic Risk Signature (MyPRS) (Cleveland Clinic Laboratories)	81599	C90.00-C90.02	15, 28
<b>Oncology: Test-Specific Not Covered Algorithmic Tests</b>				
<a href="#">Oncology: Test-Specific Not Covered Algorithmic Tests</a>	Onco4D (Animated Dynamics, Inc.)	0083U		
	BBDRisk Dx (Silbiotech)	0067U		
	PreciseDx Breast Cancer Test (PreciseDx)	0220U		
	Lymph3Cx Lymphoma Molecular Subtyping Assay (Mayo Clinic Laboratories)	0120U		
	LC-MS/MS Targeted Proteomic Assay (OncoOmicDx laboratory)	0174U		

## OTHER RELATED POLICIES

This policy document provides coverage criteria for tests that determine the risk for or the prognosis for cancer. For other oncology related testing, please refer to:

- **Oncology: Molecular Analysis of Solid Tumors and Hematologic Malignancies** for criteria related to DNA testing of a solid tumor or a blood cancer.
- **Genetic Testing: Hereditary Cancer Susceptibility Syndromes** for criteria related to genetic testing to determine if an individual has an inherited cancer susceptibility syndrome.
- **Oncology: Cancer Screening** for criteria related to the use of non-invasive fecal, urine or blood tests for screening for cancer.
- **Oncology: Circulating Tumor DNA and Circulating Tumor Cells (Liquid Biopsy)** for criteria related to circulating tumor DNA (ctDNA) or circulating tumor cell

testing performed on peripheral blood for cancer diagnosis, management and surveillance.

- **Genetic Testing: General Approach to Genetic Testing** for coverage criteria related to algorithmic testing in oncology that is not specifically discussed in this or another non-general policy.

[back to top](#)

## COVERAGE CRITERIA

### BREAST CANCER

#### Breast Cancer Treatment and Prognostic Algorithmic Tests

- I. The use of the breast cancer treatment and prognostic algorithmic test Oncotype DX Breast Recurrence Score (81519, S3854) is considered **medically necessary** in all patients, regardless of gender, when:
  - A. The member has primary breast cancer that is [ductal/NST](#), lobular, mixed or micropapillary, **AND**
  - B. The member's tumor is hormone receptor-positive (estrogen receptor-positive or progesterone receptor-positive), **AND**
  - C. The member's tumor is human epidermal growth factor receptor 2 (HER2)-negative, **AND**
  - D. The member is considering treatment with adjuvant therapy (for example, tamoxifen, aromatase inhibitors, immunotherapy), **AND**
  - E. The member meets one of the following (regardless of menopausal status):
    1. Tumor is greater than 0.5 cm and node negative (pN0), **OR**
    2. Lymph nodes are pN1mi (2mm or smaller axillary node metastases), **OR**

3. Lymph nodes are pN1 (1-3 positive nodes), **OR**
- II. The use of the breast cancer treatment and prognostic algorithmic test Breast Cancer Index (BCI) (S3854, 81518) is considered **medically necessary** when:
    - A. The member is female, **AND**
    - B. The member has primary breast cancer that is [ductal/NST](#), lobular, mixed or micropapillary, **AND**
    - C. The member's tumor is hormone receptor-positive (estrogen receptor-positive or progesterone receptor-positive), **AND**
    - D. The member's tumor is human epidermal growth factor receptor 2 (HER2)-negative, **AND**
    - E. The member is considering treatment with adjuvant therapy (for example, tamoxifen, aromatase inhibitors, immunotherapy), **AND**
      1. The member meets one of the following based on menopausal status:
        - a) The member is premenopausal and meets one of the following:
          - (1) Tumor is greater than 0.5 cm and node negative (pN0), **OR**
          - (2) Lymph nodes are pN1mi (2mm or smaller axillary node metastases), **OR**
          - (3) Lymph nodes are pN1 (1-3 positive nodes), **OR**
        - b) The member is postmenopausal and meets one of the following:
          - (1) Tumor is greater than 0.5 cm, **OR**
          - (2) Lymph nodes are pN1mi (2mm or smaller axillary node metastasis), **OR**



(3) Lymph nodes are pN1 (1-3 positive nodes).

- III. The use of the breast cancer treatment and prognostic algorithmic test Breast Cancer Index (BCI) (81518, S3854) in men with breast cancer is considered **investigational**.
- IV. The use of a breast cancer treatment and prognostic algorithmic test (i.e., Oncotype DX, Breast Recurrence Score, or Breast Cancer Index) (81518, 81519, S3854) is considered **investigational** for all other indications.

[back to top](#)

## Breast Cancer Prognostic Algorithmic Tests

- I. The use of a breast cancer prognostic algorithmic test (i.e., Endopredict, Prosigna, Mammaprint) (S3854, 81520, 81521, 81522, 81523) is considered **medically necessary** when:
  - A. The member is female, **AND**
  - B. The member has primary breast cancer that is [ductal/NST](#), lobular, mixed or micropapillary, **AND**
  - C. The member's tumor is hormone receptor-positive (estrogen receptor-positive or progesterone receptor-positive), **AND**
  - D. The member's tumor is human epidermal growth factor receptor 2 (HER2)-negative, **AND**
  - E. The member is considering treatment with adjuvant therapy (for example, tamoxifen, aromatase inhibitors, immunotherapy), **AND**
  - F. The member meets one of the following based on menopausal status:
    - 1. The member is premenopausal and meets one of the following:
      - a) Tumor is greater than 0.5 cm and node negative (pN0), **OR**

- b) Lymph nodes are pN1mi (2mm or smaller axillary node metastases), **OR**
    - c) Lymph nodes are pN1 (1-3 positive nodes), **OR**
  - 2. The member is postmenopausal and meets one of the following:
    - a) Tumor is greater than 0.5 cm, **OR**
    - b) Lymph nodes are pN1mi (2mm or smaller axillary node metastasis), **OR**
    - c) Lymph nodes are pN1 (1-3 positive nodes).
- II. The use of a breast cancer prognostic algorithmic test (i.e., Endopredict, Prosigna, Mammaprint) (S3854, 81520, 81521, 81522, 81523) in men with breast cancer is considered **investigational**.
- III. The use of a breast cancer prognostic algorithmic test (i.e., Endopredict, Prosigna, Mammaprint ) (S3854, 81520, 81521, 81522, 81523) is considered **investigational** for all other indications.

[back to top](#)

## Gene Expression Profiling Breast Cancer Subtyping Tests

- I. Gene expression profiling breast cancer subtyping tests (e.g., BluePrint) (81599, S3854, 0153U) are considered **investigational**.

[back to top](#)

## Breast DCIS Prognostic Algorithmic Tests

- I. Breast DCIS prognostic algorithmic tests (0045U) are considered **investigational**.

[back to top](#)

## COLORECTAL CANCER

### Colorectal Cancer Prognostic Algorithmic Tests

- I. Colorectal cancer prognostic algorithmic tests (81525, 0069U) are considered **investigational**.

[back to top](#)

## PROSTATE CANCER

### Prostate Cancer Treatment and Prognostic Algorithmic Tests

- I. The use of a prostate cancer treatment and prognostic algorithmic test (i.e., Oncotype DX Prostate (0047U), Prolaris (81541)) is considered **medically necessary** when:
  - A. The member has a life expectancy of 10 years or more, **AND**
  - B. The member has any of the following:
    1. [Low-risk prostate cancer](#), **OR**
    2. [Favorable intermediate prostate cancer](#), **OR**
    3. [Unfavorable intermediate prostate cancer](#), **OR**
    4. [High-risk prostate cancer](#).
- II. The use of the prostate cancer treatment and prognostic algorithmic test Decipher assay (81542) is considered **medically necessary** when:
  - A. The member meets the following:
    1. The member has a life expectancy of 10 years or more, **AND**
    2. The member has any of the following:
      - a) [Low-risk prostate cancer](#), **OR**

- b) [Favorable intermediate prostate cancer](#), **OR**
- c) [Unfavorable intermediate prostate cancer](#), **OR**
- d) [High-risk prostate cancer](#), **OR**

B. The member meets the following:

1. The test is being used to inform adjuvant treatment and counseling for risk stratification, **AND**
2. Adverse features were found post-radical prostatectomy, including but not limited to PSA resistance/recurrence.

III. The use of a prostate cancer treatment and prognostic algorithmic test (0047U, 81541, 81542) is considered **investigational** for all other indications.

[back to top](#)

## Prostate Cancer Risk Assessment Algorithmic Tests

- I. Prostate cancer risk assessment algorithmic tests (81539, 84153, 84154, 86316, 0339U, 0005U, 0359U) are considered **investigational**.

[back to top](#)

## Prostate Cancer Diagnostic Algorithmic Tests

- I. Prostate cancer diagnostic algorithmic tests (81551, 0113U) are considered **investigational**.

[back to top](#)

## THYROID CANCER

### Thyroid Cancer Diagnostic Algorithmic Tests

- I. The use of a thyroid cancer diagnostic algorithmic test (0026U, 0018U, 0204U, 0245U, 81546) in fine needle aspirates of thyroid nodules is considered **medically necessary** when:
  - A. The fine needle aspirate showed [indeterminate cytologic findings](#), **AND**
  - B. Clinical and/or radiologic findings of the thyroid nodules are indeterminate of malignancy, **AND**
  - C. The result of the test would affect surgical decision making.
- II. The use of a thyroid cancer diagnostic algorithmic test (0026U, 0018U, 0204U, 0245U, 81546) in fine needle aspirates of thyroid nodules is considered **investigational** for all other indications.

[back to top](#)

## UVEAL MELANOMA

### Uveal Melanoma Prognostic Algorithmic Tests

- I. The use of a uveal melanoma prognostic algorithmic test (81552) is considered **medically necessary** when:
  - A. The member has primary, localized uveal melanoma.
- II. The use of a uveal melanoma prognostic algorithmic test (81552) is considered **investigational** for all other indications.

[back to top](#)

## CUTANEOUS MELANOMA

### Cutaneous Melanoma Prognostic Algorithmic Tests

- I. Cutaneous melanoma prognostic algorithmic tests (81529) are considered **investigational**.

[back to top](#)

### Cutaneous Melanoma Diagnostic Algorithmic Tests

- I. Cutaneous melanoma diagnostic algorithmic tests (0090U) are considered **medically necessary** when:
  - A. The member has a melanocytic neoplasm that is diagnostically uncertain or equivocal after histopathology.
- II. Cutaneous melanoma diagnostic algorithmic tests (0090U) are considered **investigational** for all other indications, including:
  - A. A melanocytic neoplasm that has pathology definitive for melanoma, desmoplastic melanoma, or sclerosing nevus.

[back to top](#)

### Cutaneous Melanoma Risk Assessment Algorithmic Tests

- I. Cutaneous melanoma risk assessment algorithmic tests (0089U) are considered **investigational**.

[back to top](#)

## OVARIAN CANCER

### Ovarian Cancer Diagnostic Algorithmic Tests

- I. Ovarian cancer diagnostic algorithmic tests (i.e., OVA1, Overa, ROMA, and OvaWatch) (0003U, 81500, 81503, 0375U) are considered **investigational** for all indications, including but not limited to:
  - A. Preoperative evaluation of adnexal masses to triage for malignancy
  - B. Screening for ovarian cancer
  - C. Selecting patients for surgery for an adnexal mass
  - D. Evaluation of patients with clinical or radiologic evidence of malignancy
  - E. Evaluation of patients with nonspecific signs or symptoms suggesting possible malignancy
  - F. Postoperative testing and monitoring to assess surgical outcome and/or to detect recurrent malignant disease following treatment

[back to top](#)

### Ovarian Cancer Treatment Algorithmic Tests

- I. Ovarian cancer treatment algorithmic tests (0172U) are considered **medically necessary** when:
  - A. The member has a diagnosis of ovarian cancer, **AND**
  - B. The member is being considered for PARP inhibitor therapy.
- II. Ovarian cancer treatment algorithmic tests (0172U) are considered **investigational** for all other indications.

[back to top](#)

## GYNECOLOGIC CANCER

### Gynecologic Cancer Treatment Algorithmic Tests

- I. Gynecologic cancer treatment algorithmic tests (81535, 81536) in the assessment of gynecological cancers are considered **investigational**.

[back to top](#)

## LUNG CANCER

### Lung Cancer Diagnostic Algorithmic Tests

- I. Lung cancer diagnostic algorithmic tests (0080U, 0092U, 81599) are considered **investigational**, including for members with undiagnosed pulmonary nodules.

[back to top](#)

### Lung Cancer Treatment Algorithmic Tests

- I. Lung cancer treatment algorithmic tests (81538, 0288U) are considered **investigational**.

[back to top](#)

## BLADDER AND URINARY TRACT CANCER

### Bladder Cancer Diagnostic and Recurrence Algorithmic Tests

- I. Bladder cancer diagnostic and recurrence algorithmic tests (0012M, 0013M, 0363U,, 0375U, 0376U, 0377U), which are performed on urine, are considered **investigational**.

[back to top](#)



## Urinary Tract Cancer Recurrence Algorithmic Tests

- I. Urinary tract cancer recurrence algorithmic tests (86386) which are typically performed on urine are considered **investigational**.

[back to top](#)

## PANCREATIC CANCER

### Pancreatic Cyst Risk Assessment Algorithmic Tests

- I. Pancreatic cyst risk assessment algorithmic tests (0313U, 81479) are considered **investigational**.

[back to top](#)

## CANCER OF UNKNOWN PRIMARY

### Cancer of Unknown Primary Gene Expression Profiling Tests

- I. The use of a cancer of unknown primary gene expression profiling test (81540) to evaluate the site of origin of a tumor of unknown primary, or to distinguish a primary from a metastatic tumor is considered **investigational**.

[back to top](#)

## POLYGENIC RISK SCORE TESTS

### Breast Cancer Polygenic Risk Score Tests

- I. The use of a breast cancer polygenic risk score test (81599) is considered **investigational**.

[back to top](#)

## Multiple Myeloma Polygenic Risk Score Tests

- I. The use of a multiple myeloma polygenic risk score test (81599) is considered **investigational**.

[back to top](#)

## ONCOLOGY: TEST-SPECIFIC NOT COVERED ALGORITHMIC TESTS

- I. The use of these specific oncology algorithmic tests are considered **investigational**:
  - A. BBDRisk Dx (0067U)
  - B. Onco4D (0083U)
  - C. Lymph3Cx Lymphoma Molecular Subtyping Assay (0120U)
  - D. PreciseDx™ Breast Cancer Test (0220U)
  - E. LC-MS/MS Targeted Proteomic Assay (OncoOmicDx laboratory) (0174U)

[back to top](#)

## CLINICAL CONSIDERATIONS

The Oncotype DX, EndoPredict, Breast Cancer Index, MammaPrint, and Prosigna assays should only be ordered on a tissue specimen obtained during surgical removal of the tumor and after subsequent pathology examination of the tumor has been completed and determined to meet the above criteria (ie, the test should not be ordered on a preliminary core biopsy). The test should be ordered in the context of a physician-patient

discussion regarding risk preferences when the test result will aid in making decisions regarding chemotherapy.

For patients who otherwise meet the criteria for gene expression profiling for breast cancer but who have multiple ipsilateral primary tumors, a specimen from the tumor with the most aggressive histologic characteristics should be submitted for testing. It is not necessary to test each tumor; treatment is based on the most aggressive lesion.

[back to top](#)

## DEFINITIONS

1. **Ductal/NST breast cancer** is ductal cancer that is no special type (NST), meaning the cancer cells have no features that class them as a special type of breast cancer when examined by microscope.
2. **Thyroid nodules with indeterminate findings** include Bethesda diagnostic category III (atypia/follicular lesion of undetermined significance) or Bethesda diagnostic category IV (follicular neoplasm/suspicion for a follicular neoplasm)
3. **Somatic** mutations can occur in any of the cells of the body except the germ cells (sperm and egg) and therefore are not passed on to children. These alterations can (but do not always) cause cancer or other diseases.
4. **Adjuvant** therapy refers to medication (such as chemotherapy or endocrine therapy) given after the surgical removal of a cancerous tumor.
5. **Prostate cancer pathology risk stratification** is described in detail in the NCCN Prostate Cancer 1.2023 guidelines (p. PROS-2).

[back to top](#)

## BACKGROUND AND RATIONALE

### BREAST CANCER

### *National Comprehensive Cancer Network (NCCN)*

NCCN Guidelines for Breast Cancer (2.2023) makes recommendations for gene expression testing when considering adjuvant systemic therapy based on characteristics of the patient and the breast cancer. These characteristics include the patient's sex, menopause status, the TNM staging of the tumor, the expression of hormone receptors, HER2 status, and how the test will be used (i.e., prognosis alone, or prognosis and treatment decisions).

## **Breast Cancer Treatment and Prognostic Algorithmic Tests**

### Oncotype DX

Oncotype DX for breast cancer is a 21-gene expression assay. NCCN guidelines for Breast Cancer (2.2023) strongly recommends consideration of the 21-gene expression assay for both prognosis and treatment decisions in the following patients:

- Patients of either sex (p. BINV-J 1 of 2)
- Evidence level 1: Postmenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is pT1–3, and at least 0.5cm, with pN1mi (2 mm or smaller axillary node metastases) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-6, BINV-N 1 of 5, BINV-N 2 of 5)
- Evidence level 1: Premenopausal patient with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN0. Tumor must be HR positive, HER2 negative. (p. BINV-7, BINV-N 1 of 5, BINV-N 2 of 5)
- Evidence level 2A: Premenopausal patient with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN1mi (2 mm or smaller axillary node metastasis) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-8, BINV-N 1 of 5, BINV-N 2 of 5)

### Breast Cancer Index (BCI)

The BCI is recommended by NCCN (Breast Cancer, 2.2023) for both indications of prognosis as well as predicting treatment for extended adjuvant endocrine therapy. Appropriate patients for this test are:

- Patients who are female (p. BINV-J 1 of 2)

- Evidence level 2A: Postmenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is pT1–3, and 0.5cm or larger, with pN1mi (2 mm or smaller axillary node metastases) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-6, BINV-N 1 of 5, BINV-N 4 of 5)
- Evidence level 2A: Premenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN0. Tumor must be HR positive, HER2 negative. (p. BINV-7, BINV-N 1 of 5, BINV-N 4 of 5)
- Evidence level 2A: Premenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN1mi (2 mm or smaller axillary node metastasis) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-8, BINV-N 1 of 5, BINV-N 4 of 5)

### **Breast Cancer Prognostic Algorithmic Tests**

While Oncotype DX for Breast Recurrence Score is preferred by NCCN (Breast Cancer, 2.2023), other tests may be considered for prognosis/recurrence risk without treatment guidelines for patients who have hormone receptor-positive breast cancer. These tests include Endopredict and Prosignia (evidence level category 2A) and Mammaprint (evidence level category 1), which are appropriate for the following patients:

- Patients who are female (p. BINV-J 1 of 2)
- Evidence level 2A: Postmenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is pT1–3, and 0.5cm or larger, with pN1mi (2 mm or smaller axillary node metastases) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-6, BINV-N 1 of 5, BINV-N 3 of 5)
- Evidence level 2A: Premenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN0. Tumor must be HR positive, HER2 negative. (p. BINV-7, BINV-N 1 of 5, BINV-N 3 of 5)
- Evidence level 2A: Premenopausal patients with a ductal/NST, lobular, mixed, or micropapillary tumor that is at least 0.5cm and pN1mi (2 mm or smaller axillary node metastasis) or pN1 (1–3 positive nodes). Tumor must be HR positive, HER2 negative. (p. BINV-8, BINV-N 1 of 5, BINV-N 3 of 5)

### **Gene Expression Profiling Breast Cancer Subtyping Tests**

### *National Comprehensive Cancer Network (NCCN)*

NCCN Breast Cancer guidelines (2.2023) do not reference gene expression profiling tests ( i.e., Blueprint) for the purpose of subtyping breast cancer to provide information for clinical decision-making.

### **Breast DCIS Prognostic Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

NCCN Breast Cancer guidelines (2.2023) do not reference DCIS prognostic algorithmic tests as part of the clinical work-up for DCIS.

## **COLORECTAL CANCER**

### **Colorectal Cancer Prognostic Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Colon Cancer (3.2022) state that there is currently insufficient data to recommend multigene panels to assist in making clinical decisions about adjuvant therapy (p. COL-3).

## **PROSTATE CANCER**

### **Prostate Cancer Treatment and Prognostic Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

Current NCCN guidelines for Prostate Cancer (1.2023) support the consideration of gene expression profiling (specifically Decipher, Oncotype DX Prostate, and Prolaris) for prognosis and management in men with low, favorable intermediate, unfavorable intermediate, or high-risk disease, and if the patient is expected to live 10 years or longer. (p. PROS-D 2 of 4)

#### *American Society of Clinical Oncology (ASCO)*

ASCO (2020) issued a guideline for the use of molecular biomarkers in localized prostate cancer that included the following summary of recommendations:

“Tissue-based molecular biomarkers (evaluating the sample with the highest volume of the highest Gleason pattern) may improve risk stratification when added to standard clinical parameters, but the Expert Panel endorses their use only in situations in which the assay results, when considered as a whole with routine clinical factors, are likely to affect a clinical decision. These assays are not recommended for routine use as they have not been prospectively tested or shown to improve long-term outcomes—for example, quality of life, need for treatment, or survival.” (p. 1474)

## **Prostate Cancer Risk Assessment Algorithm Tests**

### *American Urological Association*

The American Urological Association (Carter et al, 2013; confirmed 2018) published guidelines on the early detection of prostate cancer and concluded that the literature supporting the use of genetic and protein biomarkers for prostate cancer screening and risk assessment provides little evidence for routine use at this time (p. 5). However, the guidelines did recognize that multiple approaches subsequent to a PSA test (e.g., urinary and serum biomarkers, imaging, risk calculators) are available for identifying men more likely to harbor prostate cancer and/or one with an aggressive phenotype. The use of such tools can be considered in men with a suspicious PSA level to inform prostate biopsy decisions (p. 17).

### *American Urological Association and Society of Abdominal Radiology*

The American Urological Association and the Society of Abdominal Radiology (Rosenkrantz et al, 2016) published joint guidelines on prostate magnetic resonance imaging and magnetic resonance imaging-targeted biopsy. The associations commented that there may be value in using genetic and protein biomarkers for prostate cancer risk in patients warranting repeat biopsy; however, further research is needed to fully assess the utility (p. 2)

### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Prostate Cancer early detection (1.2023) state that biomarkers meant to improve the specificity of detection of prostate cancer are not yet mandated as a first-line screening test in conjunction with serum prostate-specific antigen (PSA) (p. PROSD-3).

### **Prostate Cancer Diagnostic Algorithmic Tests**

*American Urological Association, American Society for Radiation Oncology, and Society of Urological Oncology*

The American Urological Association, American Society for Radiation Oncology, and the Society of Urologic Oncology (Sanda et al, Part 1 2017, Part 2 2018) published joint guidelines on the management of clinically localized prostate cancer which state that among most low-risk localized prostate cancer patients, genomic biomarkers have not demonstrated a clear role in the selection of active surveillance (Part 1, p. 686) or in the follow-up of patients on active surveillance. (Part 2, p. 991)

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Prostate Cancer early detection (1.2023) state that biomarkers meant to improve the specificity of detection of prostate cancer are not yet mandated as a first-line screening test in conjunction with serum prostate-specific antigen (PSA) (page PROSD-3).

## **THYROID CANCER**

### **Thyroid Cancer Diagnostic Algorithmic Tests**

*American Thyroid Association*

The American Thyroid Association (2016) updated its guidelines on the management of thyroid nodules and differentiated thyroid cancer in adults. These guidelines made the following statements on molecular diagnostics in thyroid nodules: “For nodules with AUS/FLUS [atypia of undetermined significance/follicular lesion of undetermined significance]... molecular testing may be used to supplement malignancy risk assessment in lieu or proceeding directly with either surveillance or diagnostic surgery.” (p. 21)



### *National Comprehensive Cancer Network (NCCN)*

Current NCCN Guidelines for Thyroid Carcinoma (3.2022) state that clinicians can consider molecular diagnostics on fine needle aspirate (FNA) results of thyroid nodules which are classified as Bethesda III or Bethesda IV if there is not high clinical and/or radiographic suspicion of malignancy. (p. THRY-1)

### *American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi*

The American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi (2016) updated their joint guidelines on molecular testing for cytologically indeterminate thyroid nodules and endorsed the following:

- *TERT* mutational analysis may improve the diagnostic sensitivity of molecular testing on cytologic samples. (p. 32)
- There is insufficient evidence to recommend either in favor of or against the use of gene expression classifiers for cytologically indeterminate nodules. (p. 10)
- With the exception of mutations such as BRAF V600E, there is insufficient evidence to recommend in favor of or against the use of mutation testing to determine the extent of surgery. (p. 10)

## **UVEAL MELANOMA**

### **Uveal Melanoma Prognostic Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Uveal Melanoma (2.2022) support gene expression profiling and chromosome analysis in all patients with uveal melanoma and further state that molecular testing for prognostication is preferred over cytology alone. (p. MS-6)

## **CUTANEOUS MELANOMA**

### **Cutaneous Melanoma Prognostic Algorithmic Tests**

### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Cutaneous Melanoma (1.2023) recognize that the use of gene expression profiling as an emerging technology to differentiate melanomas at low versus high risk for metastasis, to clarify indeterminate melanocytic neoplasms following histopathology, and to classify cutaneous melanoma into separate categories based on metastasis; however, currently there is insufficient data to recommend the use of gene expression profiling for cutaneous melanoma as the impact of these tests has not been established. (p. ME-C 1 of 8)

### *American Academy of Dermatology*

The American Academy of Dermatology (2019) published guidelines of care for the management of primary cutaneous melanoma. The guidelines state the following regarding GEP tests:

- There is insufficient evidence of benefit to recommend routine use of currently available prognostic molecular tests, including GEP, for prognosis of CM. (page 219)
- Routine molecular testing, including GEP, for prognostication is discouraged until better use criteria are defined. The application of molecular information for clinical management is not recommended. (p. 219)

## **Cutaneous Melanoma Diagnostic Algorithmic Tests**

### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Cutaneous Melanoma (1.2023) indicate that gene expression profiling is an acceptable test for diagnosing indeterminate melanocytic neoplasms by histopathology, along with immunohistochemistry (IHC), comparative genomic hybridization (CGH), fluorescence in situ hybridization (FISH), single-nucleotide polymorphism (SNP) array, and next-generation sequencing (NGS). These tests may lead to a definitive diagnosis and guide therapy in cases that are diagnostically uncertain or controversial by histopathology. (p. ME-C 1 of 8).

### *American Academy of Dermatology*

The American Academy of Dermatology (Swetter, 2019) published guidelines of care for the management of primary cutaneous melanoma. The guidelines state the following regarding GEP tests:

- Diagnostic molecular techniques are still largely investigative and may be appropriate as ancillary tests in equivocal melanocytic neoplasms, but they are not recommended for routine diagnostic use in CM. These include comparative genomic hybridization (CGH), fluorescence in situ hybridization (FISH), gene expression profiling (GEP), and (potentially) next-generation sequencing. (page 219)
- Ancillary diagnostic molecular techniques (eg, CGH, FISH, GEP) may be used for equivocal melanocytic neoplasms. (p. 219)

#### *American Society of Dermatopathology*

The American Academy of Dermatopathology (AUC Committee Members, 2022) published conditions where a 23 gene qRT-PCR test (MyPath Melanoma) was determined by a review of published evidence to be “majority usually appropriate.” These include the differential diagnosis of nevus versus melanoma in fully sampled histopathologically ambiguous tumors, partially sampled nevus versus melanoma in adults, nevus versus nevoid melanoma, and nevus versus melanoma in cosmetically sensitive sites and special sites in pediatric patients. These recommendations specifically exclude scenarios where pathology is definitive for melanoma or for distinction between incompletely sampled sclerosing (desmoplastic) nevus versus desmoplastic melanoma. (p. 237-8)

### **Cutaneous Melanoma Risk Assessment Algorithmic Tests**

#### *Concert Genetics*

This review focused on peer-reviewed, published evidence of the clinical utility of DermTech PLA through May, 2022. PubMed and ECRI Guidelines Trust searches were performed. Search terms included pigmented lesion assay, DermTech, 0089U, PRAME, LINC00518, cutaneous melanoma risk. References were also identified from the performing laboratory’s website. A total of 110 abstracts from these sources were reviewed, and 30 full text publications were evaluated. At the present time, the DermTech Pigmented Lesion Assay has not been adequately shown in peer-reviewed

publications to effectively result in improved health outcomes compared to the current standard of care.

## **OVARIAN CANCER**

### **Ovarian Cancer Diagnostic Algorithmic Tests**

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Ovarian Cancer, Fallopian Tube Cancer, and Primary Peritoneal Cancer (1.2023) recognize the use of biomarker analysis for risk assessment for ovarian cancer in women with a pelvic mass as an emerging technology; however, the NCCN panel of experts currently does not recommend these biomarker tests for clinical use. (p. MS-10 and p. MS-11)

### **Ovarian Cancer Treatment Algorithmic Tests**

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Ovarian Cancer, Fallopian Tube Cancer, and Primary Peritoneal Cancer (1.2023) recommend genetic risk evaluation, and germline and somatic testing if not previously done, including *BRCA1/2* to inform maintenance therapy for patients with ovarian, fallopian tube, or primary peritoneal cancer. If a patient does not have a germline *BRCA1/2* mutation, homologous recombination status may inform on the benefit of PARP inhibitor therapy. (p. OV-1)

*American Society of Clinical Oncology (ASCO)*

ASCO (2020) issued a guideline for the use of PARP inhibitors in the management of ovarian cancer, which included the following summary of recommendations:

“The guideline pertains to patients who are PARPi naïve. All patients with newly diagnosed, stage III-IV EOC (epithelial ovarian, tubal, or primary peritoneal cancer), whose disease is in complete or partial response to first-line, platinum-based chemotherapy with high-grade serous or endometrioid EOC should be offered PARPi maintenance therapy with niraparib. For patients with germline or somatic pathogenic or likely pathogenic variants in *BRCA1* (g/sBRCA1) or *BRCA2* (g/sBRCA2) genes, should be treated with olaparib. The addition of olaparib to

bevacizumab may be offered to patients with stage III-IV EOC with g/sBRCA1/2 and/or genomic instability and a partial or complete response to chemotherapy plus bevacizumab combination. Maintenance therapy (second line or more) with single-agent PARPi may be offered for patients with EOC who have not received a PARPi and have responded to platinum-based therapy regardless of BRCA mutation status. Treatment with a PARPi should be offered to patients with recurrent EOC that has not recurred within 6 months of platinum-based therapy, who have not received a PARPi and have a g/sBRCA1/2, or whose tumor demonstrates genomic instability. PARPis are not recommended for use in combination with chemotherapy, other targeted agents, or immune-oncology agents in the recurrent setting outside the context of a clinical trial. Recommendations for managing specific adverse events are presented. Data to support reuse of PARPis in any setting are needed.” (p. 3)

## **GYNECOLOGIC CANCER**

### **Gynecologic Cancer Treatment Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Ovarian c=Cancer, Fallopian Tube Cancer, and Primary Peritoneal Cancer (1.2023) state that chemosensitivity or chemoresistance assays, or other biomarker assays, are being used at some institutions, but the current level of evidence is not sufficient to replace the current standard of care of chemotherapy (p. OV-C).

NCCN guidelines for Cervical Cancer (1.2023) do not mention chemosensitivity or chemoresistance assays as part of clinical care.

NCCN guidelines for Uterine Neoplasms (1.2023) do not mention chemosensitivity or chemoresistance assays as part of clinical care.

## **LUNG CANCER**

### **Lung Cancer Treatment Algorithmic Tests**

#### *Concert Genetics*

This review focused on peer-reviewed, published evidence of the clinical utility of VeriStrat® through June 2022. PubMed and ECRI Guidelines Trust searches were performed. Search terms included VeriStrat®, proteomic non-small cell lung cancer, prognosis, and survival. References were also identified from the performing laboratory's website. At the present time, the VeriStrat® test has not been adequately shown in peer-reviewed publications to effectively result in improved health outcomes compared to the current standard of care.

## **Lung Cancer Diagnostic Algorithmic Tests**

### *Concert Genetics*

This review focused on peer-reviewed, published evidence of the clinical utility of NodifyXL2, Percepta Bronchial Genomic Classifier, and Reveal Lung Nodule Characterization through June 2022. PubMed and ECRI Guidelines Trust searches were performed. Search terms included Nodify, Percepta, lung nodule, plasma-protein and multiplex. References were also identified from the performing laboratory's website. At the present time, NodifyXL2, Percepta Bronchial Genomic Classifier, and Reveal Lung Nodule Characterization have not been adequately shown in peer-reviewed publications to effectively result in improved health outcomes compared to the current standard of care.

## **BLADDER AND URINARY TRACT CANCER**

### **Bladder Cancer Diagnostic and Recurrence Algorithmic Tests**

#### *National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Bladder Cancer (1.2023) support consideration for urinary urothelial tumor markers for high-risk patients with non-muscle-invasive bladder cancer (category 2B recommendation) (p. BL-E 2 of 6). Further discussion in these guidelines acknowledge that it is unclear if this type of testing offers information that is clinically useful for detecting or managing these tumors, hence the weaker recommendation of 2B by the panel (p. MS-13)

#### *American Urological Association and Society of Urologic Oncology*

The American Urological Association and Society of Urologic Oncology (Chang et al, 2016) addressed the diagnosis and treatment of non-muscle-invasive bladder cancer, based on a systematic review and includes the following statements on the use of urine markers after the diagnosis of bladder cancer:

- Urinary biomarker analysis should not replace cystoscopic evaluation in the surveillance of NMIBC [non-muscle invasive bladder cancer]. (Strong Recommendation; Evidence Strength: Grade B)
- Urinary biomarker analysis or cytology should not routinely be used during surveillance in a patient with a history of low-risk cancer and a normal cystoscopy (Expert Opinion) (p. 1024 and 1025)

Note: “Evidence Strength B” describes a recommendation of moderate certainty. “Expert Opinion” is defined in this guideline as “A statement, achieved by consensus of the Panel, that is based on members’ clinical training, experience, knowledge, and judgment for which there is no evidence.” (p. 1022)

### **Urinary Tract Cancer Recurrence Algorithmic Tests**

Current NCCN guidelines on Bladder Cancer (1.2023) does not include a recommendation for algorithmic-based screening for urinary tract cancer. .

## **PANCREATIC CANCER**

### **Pancreatic Cyst Risk Assessment Algorithmic Tests**

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Pancreatic Adenocarcinoma (2.2022) discuss the use of endoscopic ultrasound to follow patients with pancreatic cysts and after the removal, citing that the risk of malignancy in mucinous cystic neoplasms is less than 15%. (p. MS-6, MS-10) The guidelines do not include recommendation or discussion for the use of molecular analysis of pancreatic cysts to stratify risk of cancer.

*American College of Gastroenterology*

The American College of Gastroenterology (2018) published guidelines for the diagnosis and management of pancreatic cysts, which included the following:

“A number of DNA, RNA, protein, and metabolomic markers have been evaluated in cyst fluid. The majority of these are still early in development and not yet ready for translation into clinical practice. However, analysis of DNA mutations in cyst fluid has shown promise in identifying IPMNs [intraductal papillary mucinous neoplasms] and MCNs [mucinous cystic neoplasms].” (p. 471)

## **CANCER OF UNKNOWN PRIMARY**

### **Cancer of Unknown Primary Gene Expression Profiling Tests**

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for Occult ]Primary (Cancer of Unknown Primary) (3.2023) state that gene sequencing to predict tissue of origin is not recommended (p. OCC-1).

## **POLYGENIC RISK SCORE TESTS**

### **Breast Cancer Polygenic Risk Score Tests**

*National Comprehensive Cancer Network (NCCN)*

NCCN guidelines for genetic/familial high-risk assessment for breast, ovarian, and pancreatic cancers (3.2023) speak broadly about the use of polygenic risk scores, stating that there are currently significant limitations to this type of testing, and it should not be used for clinical management at this time outside of the context of a clinical trial (p. EVAL-A).

### **Multiple Myeloma Polygenic Risk Score Tests**

*National Comprehensive Cancer Network (NCCN)*



NCCN guidelines for Multiple Myeloma (3.2023) do not mention the use of polygenic risk score as part of clinical management for multiple myeloma.

NCCN guidelines for genetic/familial high-risk assessment for breast, ovarian, and pancreatic cancers (3.2023) speak broadly about the use of polygenic risk scores, stating that there are currently significant limitations to this type of testing, and it should not be used for clinical management at this time outside of the context of a clinical trial (p. EVAL-A).

[back to top](#)

## REFERENCES

1. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 2.2023.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf)
2. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. Version 3.2022.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/colon.pdf](https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf)
3. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer. Version 1.2023.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/prostate.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf)
4. Carter HB, Albertsen PC, Barry MJ, et al. Early Detection of Prostate Cancer: AUA Guideline. J Urol. 2013;190(2):419-426. doi:10.1016/j.juro.2013.04.119.
5. Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options. J Urol. 2018;199(3):683-690. doi:10.1016/j.juro.2017.11.095.
6. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Thyroid Cancer. Version 3.2022.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/thyroid.pdf](https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf)
7. Gharib H, Papini E, Garber JR, et al. American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi Medical Guidelines for Clinical Practice for the Diagnosis and Management of Thyroid Nodules--2016 Update. Endocr Pract. 2016;22(5):622-639. doi:10.4158/EP161208.GL.

8. Haugen BR, Alexander EK, Bible KC, et al. American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. 2016;26(1):1-133. doi:10.1089/thy.2015.0020.
9. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Uveal Melanoma. Version 2.2022. [https://www.nccn.org/professionals/physician\\_gls/pdf/uveal.pdf](https://www.nccn.org/professionals/physician_gls/pdf/uveal.pdf)
10. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Cutaneous Melanoma. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/cutaneous\\_melanoma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf)
11. Swetter, SS, Tsao, HH, Bichakjian, CC, et al. Guidelines of care for the management of primary cutaneous melanoma. *J Am Acad Dermatol*. 2019;80(1):208-250. doi:10.1016/j.jaad.2018.08.055.
12. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Ovarian Cancer including Fallopian Tube Cancer and Primary Peritoneal Cancer. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/ovarian.pdf](https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf)
13. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Bladder Cancer. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/bladder.pdf](https://www.nccn.org/professionals/physician_gls/pdf/bladder.pdf)
14. Chang SS, Boorjian SA, Chou R, et al. Diagnosis and treatment of non-muscle invasive bladder cancer: AUA/SUO Guideline. *J Urol*. 2016;196(4):1021-1029. doi:10.1016/j.juro.2016.06.049.
15. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Genetic/Familial High Risk Assessment: Breast, Ovarian and Pancreatic. Version 3.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/genetics\\_bop.pdf](https://www.nccn.org/professionals/physician_gls/pdf/genetics_bop.pdf)
16. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Occult Primary (Cancer of Unknown Primary). Version 3.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/occult.pdf](https://www.nccn.org/professionals/physician_gls/pdf/occult.pdf)
17. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Cervical Cancer. Version 1.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/cervical.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cervical.pdf)

18. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Uterine Neoplasms. Version 1.2023.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/uterine.pdf](https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf)
19. Eggener SE, Rumble RB, Armstrong AJ, et al. Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. J Clin Oncol. 2020;38(13):1474-1494. doi:10.1200/JCO.19.02768
20. Tew WP, Lacchetti C, Ellis A, et al. PARP Inhibitors in the Management of Ovarian Cancer: ASCO Guideline. J Clin Oncol. 2020;38(30):3468-3493. doi:10.1200/JCO.20.01924
21. Elta GH, Enestvedt BK, Sauer BG, Lennon AM. ACG Clinical Guideline: Diagnosis and Management of Pancreatic Cysts. Am J Gastroenterol. 2018;113(4):464-479. doi:10.1038/ajg.2018.14
22. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Pancreatic Adenocarcinoma. Version 2.2022.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/pancreatic.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf)
23. Concert Genetics. Evidence Review for Coverage Determination for Cutaneous Melanoma Risk Assessment Algorithmic Tests. V1.2.2022
24. Concert Genetics. Evidence Review for Coverage Determination for Lung Cancer Treatment Algorithmic Tests. V1.2.2022
25. Concert Genetics. Evidence Review for Coverage Determination for Lung Cancer Diagnostic Algorithmic Tests. V1.2.2022
26. AUC Committee Members, Fung MA, Vidal CI, et al. Appropriate use criteria for ancillary diagnostic testing in dermatopathology: New recommendations for 11 tests and 220 clinical scenarios from the American Society of Dermatopathology Appropriate Use Criteria Committee. J Cutan Pathol. 2022;49(3):231-245. doi:10.1111/cup.14135
27. Rosenkrantz AB, Verma S, Choyke P, et al. Prostate Magnetic Resonance Imaging and Magnetic Resonance Imaging Targeted Biopsy in Patients with a Prior Negative Biopsy: A Consensus Statement by AUA and SAR. J Urol. 2016;196(6):1613-1618. doi:10.1016/j.juro.2016.06.079
28. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Multiple Myeloma. Version 3.2023.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf)
29. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer Early Detection. Version 1.2023.  
[https://www.nccn.org/professionals/physician\\_gls/pdf/prostate\\_detection.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate_detection.pdf)

[back to top](#)