



<b>Magellan Healthcare</b>	
<b>Clinical guidelines</b> <b>PASSIVE TREATMENT</b>	<b>Original Date:</b> November 2, 2015 <b>Page 1 of 12</b>
<b>Physical Medicine – Clinical Decision Making</b>	<b>Last Review Date:</b> June 2017
<b>Guideline Number: NIA_CG_604</b>	<b>Last Revised Date:</b> September 2017
<b>Responsible Department:</b> <b>Clinical Operations</b>	<b>Implementation Date:</b> January 2018

### **Policy Statement**

Magellan Healthcare does not support the use of multiple passive treatments for the care of musculoskeletal pain within the scope of network practitioners. Most passive treatments have similar physiological effects related to pain control and reduction of inflammation. The use of modalities with duplicative physiological effects is unnecessary and inappropriate. Multiple passive treatments have not been shown to improve or accelerate patient health outcomes.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

### **Purpose**

This policy will be used to provide medical necessity guidelines to support passive treatment services for musculoskeletal conditions in a clinical setting.

### **Scope**

Physical medicine participating network practitioners, including rendering chiropractors, physical therapists, occupational therapists, speech therapists and therapist assistants as applicable. This policy also applies to out of network practitioners as dictated by the health plan.

### **Definition**

#### Modality:

Modality is defined as any group of agents that may include thermal, acoustic, radiant, mechanical, or electrical energy to produce physiologic changes in tissues of therapeutic purposes. Modalities affect tissue at the cellular level.

#### Multiple Passive Modalities:

Multiple passive modalities are defined as the use of and/or billing of two or more physical medicine modalities each visit or during the same session to the same region.

#### Passive Modalities:

Modality that is applied by the provider or in a clinical setting and does not involve active participation by the patient. The purpose of passive modalities use is to promote pain reduction, improve function and quickly transition the patient to self-care engagement.

#### Procedure:

Procedure is a service provided to increase the functional abilities in self-care, mobility, or safety.

I. The following is a list of procedures and modalities considered to be passive treatment:

- A. Thermal and light therapy – Hot/cold (97010), diathermy (97024), microwave (97020) infrared (97026), ultraviolet (97028), ultrasound (97035), paraffin bath (97018) and whirlpool (97022).
- B. Electrical therapy – High volt, low volt, interferential current, TENS (97014 and 97032).
- C. Mechanical – mechanically assisted and often a sustained pull of the spine or limb such as traction (97012). The use of traction for low back pain, with or without sciatica, is not supported by the literature, and is therefore not considered medically necessary.
- D. Therapeutic massage and manual therapy (97124 and 97140). Manual therapy includes Active Release Technique, trigger point therapy, myofascial release, mobilization/manipulation, manual lymphatic drainage, and manual traction.

II. **Appropriate use of passive treatment:**

Passive treatment modalities may be utilized in the initial acute stage of a condition for pain control, reduction of inflammation, or reduction of muscle spasm. As a condition progresses, passive care should be replaced by active treatment modalities such as therapeutic exercise. Insufficient evidence exists to support the continued use of passive treatment as a means for improved clinical outcomes.

Passive modalities are considered to be clinically appropriate and/or necessary in the conservative management of neuromusculoskeletal conditions when:

- There are no contraindications to the intervention
- Self-administration is implausible or places the patient at risk of harm
- Used primarily during the initial period of an episode of treatment
- Used to support an active care approach (i.e., therapeutic exercise)
- Used for a particular condition for which there is an evidence-basis of significant benefit

Passive modalities are considered NOT to be clinically appropriate and/or necessary when:

- Patient safety is jeopardized by the application of the modality
- The modality can be safely self-administered
- Used during a course of treatment, which continues beyond the initial period
- Used as the primary or sole therapy
- Greater than one passive modality is used involving the same body region(s)
- Used largely for the comfort and convenience of the patient
- Used as part of the routine office protocol

### **III. Exclusions:**

The use of chiropractic manipulation (98940-98943) is not considered a duplication of service or physiological effect when used in conjunction with passive treatment modalities, except for the following:

The National Correct Coding Initiative (NCCI) edits require that the manual therapy techniques be performed in a separate anatomic site than the chiropractic adjustments in order to be reimbursed separately.

#### **Additional Information:**

The preponderance of evidence appears to support either a lack of efficacy or insufficient data to make a judgment on benefit for the modalities evaluated. When a positive outcome was described, the reported treatment effects were modest. Similarly, the duration of treatment effectiveness was typically reported as short (2 weeks to 2 months). Most international guidelines recommend these interventions should only be used reservedly based upon individual circumstances, and not as a principle component of a treatment regime.

The use of passive modalities in the treatment of neuromusculoskeletal conditions presents the inherent risk of promoting passive dependence. It is the responsibility of the treating practitioner to judiciously apply passive modalities and encourage active patient participation in the treatment plan. Passive treatment is generally viewed as appropriate when used for a short period of time and in conjunction with an active care.

A review on non-pharmacological therapies for acute and chronic LBP by the American Pain Society and the American College of Physicians (Chou et al, 2007) concluded that therapies with good evidence of moderate efficacy for chronic or sub-acute LBP are cognitive-behavioral therapy, exercise, spinal manipulation, and interdisciplinary rehabilitation. For acute LBP, the only therapy with good evidence of efficacy is superficial heat.

No high quality evidence was found to support the use of ultrasound for improving pain or quality of life in patients with non-specific chronic LBP. There is some evidence that therapeutic ultrasound has a small effect on improving low-back function in the short term, but this benefit is unlikely to be clinically important. Evidence from comparisons between other treatments and therapeutic ultrasound for chronic LBP were indeterminate and generally of low quality. There was little evidence that active therapeutic ultrasound is more effective than placebo ultrasound for treating people with pain or a range of musculoskeletal injuries or for promoting soft tissue healing. Based on low to moderate level evidence, therapeutic US does not provide any benefit compared to a placebo or advice, to laser therapy or when combined to exercise for treatment of rotator cuff tendinopathy. Ultrasound provided no additional benefit in improving pain and function in addition to exercise training in the management of knee osteoarthritis.

No trials at low risk of bias support the use of traction, stretching, or ultrasound therapy for chronic neck pain.

Overall, there was limited high quality evidence for the effectiveness of manual therapy. Most reviewed evidence was of low to moderate quality and inconsistent due to substantial methodological and clinical diversity.

No high-quality evidence was found, indicating that there is great uncertainty about the effectiveness of exercise and manual therapy for treatment of temporomandibular joint dysfunction.

For adults with nonspecific shoulder pain of variable duration, cervicothoracic spinal manipulation and mobilization in addition to usual care may improve self-perceived recovery compared to usual care alone. For adults with subacromial impingement syndrome of variable duration, neck mobilization in addition to a multimodal shoulder program of care provides no added benefit. Finally, for adults with grade I-II ankle sprains of variable duration, lower extremity mobilization in addition to home exercise and advice provides greater short-term improvements in activities and function over home exercise and advice alone.

For patients with rotator cuff tendinopathy, based on low to moderate-quality evidence, manual therapy may decrease pain; however, it is unclear whether it can improve function.

Best available evidence indicates that exercise therapy (whether land-based or water-based) is more effective than minimal control in managing pain associated with hip OA in the short term. Larger high-quality RCTs are needed to establish the effectiveness of exercise and manual therapies in the medium and long term.

Low quality evidence suggests clinically important long-term improvements in pain, function/disability, and global perceived effect when manual therapy and exercise are compared to no treatment. High quality evidence suggests greater short-term pain relief than exercise alone, but no long-term differences across multiple outcomes for (sub) acute/chronic neck pain with or without cervicogenic headache. Moderate quality evidence supports this treatment combination for pain reduction and improved quality of life over manual therapy alone for chronic neck pain; and suggests greater short-term pain reduction when compared to traditional care for acute whiplash. Evidence regarding radiculopathy was sparse.

Both stretching exercise and manual therapy considerably decreased neck pain and disability in women with non-specific neck pain. The difference in effectiveness between the 2 treatments was minor. Low-cost stretching exercises can be recommended in the first instance as an appropriate therapy intervention to relieve pain, at least in the short-term.

For the treatment of the diagnostic label Non-Specific Neck Pain strong evidence of efficacy was only found for multimodal care (manipulation/mobilization and supervised exercises).

The Cochrane Back and Neck Group reported little confidence that massage is an effective treatment for LBP. Acute, sub-acute and chronic LBP had improvements in pain outcomes with massage only in the short-term follow-up. Functional improvement was observed in participants with sub-acute and chronic LBP when compared with inactive controls, but only for the short-term follow-up.

There are insufficient data to draw firm conclusion on the clinical effect of back schools, low-level laser therapy, patient education, massage, traction, superficial heat/cold, and lumbar supports for chronic LBP.

A number of pharmacological and nonpharmacological noninvasive treatments for low back pain are associated with small to moderate, primarily short-term, effects on pain versus placebo, sham, wait list, or no treatment. Effects on function are generally smaller than effects on pain. More research is needed to understand optimal selection of treatments, effective combinations and sequencing of treatments, and effectiveness of treatments for radicular low back pain.

Guidelines on treatment of LBP from the National Collaborating Centre for Primary Care (Savigny et al, 2009) found insufficient evidence for the use of interferential stimulation in LBP and recommended against its use for that indication.

In a systematic review and meta-analysis, Fuentes et al (2010) analyzed the available information regarding the efficacy of interferential therapy in the management of musculoskeletal pain. Interferential current alone was not significantly better than placebo or other therapy at discharge or follow-up.

The effectiveness of high-voltage pulsed current treatments in humans as a means of controlling edema and post-traumatic pain has not yet been established.

Scientific evidence in peer review literature is lacking regarding the use, safety, improvement or effectiveness on health outcomes for light emitting diode (infrared) therapy.

Documentation requirements:

The treatment plan or plan of care must include the clinical rationale for each service, a description of the service, the area of the body the service will be provided, goals for each service, and a time component, if indicated.

Contraindications: The use of ultrasound therapy is contraindicated for pregnant patients or patients with malignancy.

## REFERENCES

- Alison R. Snyder, April L. Perotti, Kenneth C. Lam, and R. Curtis Bay. Influence of High-Voltage Electrical Stimulation on Edema Formation After Acute Injury: A Systematic Review. *Journal of Sport Rehabilitation*, 2010, 19, 436-451
- American College of Occupational and Environmental Medicine (ACOEM); 2nd ed. Elk Grove Village (IL); 2007. Low back disorders. Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 366
- Anne Thackeray, PT, PhD, MPH, Julie M. Fritz, PT, PhD, John D. Childs, PT, PhD, MBA, Gerard P. Brennan, PT, PhD. The Effectiveness of Mechanical Traction Among Subgroups of Patients With Low Back Pain and Leg Pain: A Randomized Trial.
- Ariel Desjardins-Charbonneau, PT, MSc, Jean-Sébastien Roy, PT, PhD, Clermont E. Dionne, OT, PhD, Pierre Frémont, MD, PhD, Joy C. MacDermid, PT, PhD, François Desmeules, PT, PhD. The Efficacy of Manual Therapy for Rotator Cuff Tendinopathy: A Systematic Review and Meta-analysis. *Journal of Orthopaedic & Sports Physical Therapy*, 2015 Volume:45 Issue:5 Pages:330–350
- B Bagheripour et al. The Efficacy of a Home-Mechanical Traction Unit for Patients With Mild to Moderate Cervical Osteoarthritis: A Pilot Study. *Med J Islam Repub Iran* 30, 386. 2016 Jun 12
- Bell J. Massage therapy helps to increase range of motion, decrease pain and assist in healing a client with low back pain and sciatica symptoms. *J Bodywork & Movement Ther* 2008; 12(3):281-289.
- Bergman S. Management of musculoskeletal pain. *Best Pract Res Clin Rheumatol*. 2007;21:153-166
- Beurskens AJ, de Vet HC, Koke AJ, et al. Efficacy of traction for nonspecific low back pain. *Spine*. 1997;22(23):2756-2762.
- Borman P, Keskin D, Ekici B, Bodur H. The efficacy of intermittent cervical traction in patients with chronic neck pain. *Clin Rheumatology* 2008; 27(10):1249-1253.
- Brosseau L, Casimiro L, Milne S, Welch V, Shea B, Tugwell P, Wells GA. Deep transverse friction massage for treating tendinitis. *Cochrane Database of Systematic Reviews* 2002, Issue 4.
- Brosseau L, Casimiro L, Welch V, Milne S, Shea B, Judd M, Wells GA, Tugwell P. Therapeutic ultrasound for treating patellofemoral pain syndrome. *Cochrane Database of Systematic Reviews* 2001, Issue 4.

Cai C, Pua Y, Lim K. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with mechanical lumbar traction. *Euro Spine J* 2009; 18(4):554-561.

Cakir S, Hepguler S, Ozturk C, Korkmaz M, Isleten B, Atamaz FC. Efficacy of therapeutic ultrasound for the management of knee osteoarthritis: a randomized, controlled, and double-blind study. *Am J Phys Med Rehabil.* 2014 May;93(5):405-12

Carey TS, Freburger JK, Holmes GM, Castel L, Darter J, Agans R, et al. A Long Way to 11 Go: Practice Patterns and Evidence in Chronic Low Back Pain Care. *Spine* 2009.

Centers for Medicare & Medicaid Services. Decision memo for transcutaneous electrical nerve stimulation for chronic low back pain (CAG-00429N). June 8, 2012. CMS: Baltimore, MD. Available at: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=256>. Accessed April 21, 2017.

Cherian JJ, Jauregui JJ, Leichter AK, et al. The effects of various physical non-operative modalities on the pain in osteoarthritis of the knee. *Bone Joint J.* 2016;98-B(1 Suppl A):89-94.

Chou R and Hoyt Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: A review of the evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline. *Annals of Internal Medicine.* 2007;147(7):492-504.

Christine Clar, Alexander Tsertsvadze, Rachel Court, Gillian Lewando Hundt, Aileen Clarke and Paul Sutcliffe. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropractic & Manual Therapies* 2014;22:12 DOI: 10.1186/2045-709X-22-12

Clarke J, van Tulder M, Blomberg S, et al. Traction for low-back pain with or without sciatica. *Cochrane Database of Systematic Reviews.* 2007 (2):CD0033010.

Clarke JA, van Tulder MW, Blomberg SE, de Vet HC, van der Heijden GJ, Bronfort G, Bouter LM. Traction for low-back pain with or without sciatica. *Cochrane Database Syst Rev.* 2007 Apr 18;(2):CD0033010.

*Cochrane Database Syst Rev.* 2014 Mar 14;(3):CD009169. doi: 10.1002/14651858.CD009169.pub2. Therapeutic ultrasound for chronic low-back pain. Ebadi S1, Henschke N, Nakhostin Ansari N, Fallah E, van Tulder MW.

Coste J, Delecoeuillerie G, Cohen de Lara, Le Parc JM, Paolaggi JB. Clinical course and prognostic factors in acute low back pain: an inception cohort study in primary care practice. *Br Med J.* 1994;308:577-80.

- Deyo RA, Walsh NE, Martin DC, et al. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N Engl J Med.* 1990;322(23):1627-34.
- E. J. Thoomes. Effectiveness of manual therapy for cervical radiculopathy, a review. *Chiropractic & Manual Therapies* 2016 24:45
- François Desmeules, Jennifer Boudreault, Jean-Sébastien Roy, Clermont Dionne, Pierre Frémont, Joy C. MacDermid. The efficacy of therapeutic ultrasound for rotator cuff tendinopathy: A systematic review and meta-analysis
- Frost H, Lamb SE, Klaber Moffett JA, et al. A fitness programme for patients with chronic low back pain: 2-year follow-up of a randomized controlled trial. *Pain.* 1998;75:273-9.
- Fuentes JP, Armijo Olivo S, Magee DJ, Gross DP. *Phys Ther.* Effectiveness of interferential current therapy in the management of musculoskeletal pain: a systematic review and meta-analysis. 2010 Sep;90(9):1219-38. doi: 10.2522/ptj.20090335. Epub 2010 Jul 22.
- Furlab A, Imamura M, Dryden T, Irvin E. Massage for low-back pain. *Cochrane Database of Systematic Reviews.* 2008 (4):CD001929.
- Furlan A, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med* 2012;953139. doi:10.1155/2012/953139
- Furlan AD, Giraldo M, Baskwill A, Irvin E, Imamura M. Massage for low-back pain. *Cochrane Database Syst Rev.* 2015 Sep;9:CD001929. PubMed #26329399. Massage for low-back pain. Updated October 12 2015
- Furlan AD, Giraldo M, Baskwill A, Irvin E, Imamura M. Massage for low-back pain. *Cochrane Database Syst Rev* 2015;9:CD001929. doi: 10.1002/14651858
- Gay R, Brault J. Evidence-informed management of chronic low back pain with traction therapy. *Spine Journal* 2008; 8(1):234-242.
- Graham N, Gross A, Goldsmith C, et al. Mechanical traction for neck pain with or without radiculopathy. *Cochrane Database of Systematic Reviews.* 2008 (3):CD006408.
- Gross AR, Hoving JL, Haines TA, et al. A Cochrane review of manipulation and mobilization for mechanical neck disorders. *Spine.* 2004;29(14):1541-1548.
- Guzman J, Haldeman S, Carroll LJ, Carragee EJ, Hurwitz EL, Peloso P, et al. Clinical practice implications of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: from concepts and findings to recommendations. *Spine* 2008;33:S199-S213.

- Hagen EM, Eriksen HR, Ursin H. Does early intervention with a light mobilization program reduce long-term sick leave for low back pain? *Spine*. 2000;25(15):1973-6.
- Haldeman S, Chapman-Smith D, Petersen DM. Frequency and duration of care. In *Guidelines for Chiropractic Quality Assurance and Practice Parameters*. Aspen 1993, Gaithersburg.
- Haraldsson B, Gross A, Myers CD, Ezzo J, Morien A, Goldsmith CH, Peloso PMJ, Brønfort G, Cervical Overview Group. Massage for mechanical neck disorders. *Cochrane Database of Systematic Reviews* 2006, Issue 3.
- Hurwitz EL, Morgenstern H, Harber P, et al. A randomized trial of medical care with and without physical therapy and chiropractic care with and without physical modalities for patients with low back pain: 6 month follow-up outcomes from the UCLA low back pain study. *Spine*. 2002;27(20):2193-204.
- Hurwitz EL, Morgenstern H, Harber P, et al. The effectiveness of physical modalities among patients with low back pain randomized to chiropractic care: findings from the UCLA low back pain study. *J Manipulative Physiol Ther*. 2002;25(1):10-20.
- Imamura M, Furlan A, Dryden T, Irvin E. Evidence-informed management of chronic low back pain with massage. *Spine Journal* 2008; 8(1): 121-133.
- Jesús Seco, MD, PhD, Francisco M. Kovacs, MN, PhD, Gerard Urrutia, MD. The efficacy, safety, effectiveness, and cost-effectiveness of ultrasound and shock wave therapies for low back pain: a systematic review. <http://dx.doi.org/10.1016/j.spinee.2011.02.002>
- Johnson M, Martinson M. Efficacy of electrical nerve stimulation for chronic musculoskeletal pain: A meta-analysis of randomized controlled trials. *Pain*. 2007;130:157-165.
- Julie M. Fritz, PT, PhD, ATC, Anne Thackeray, PT, MPH, Gerard P. Brennan, PT, PhD, FAAOMPT, John D. Childs, PT, PhD, FAAOMPT. Exercise Only, Exercise With Mechanical Traction, or Exercise With Over-Door Traction for Patients With Cervical Radiculopathy, With or Without Consideration of Status on a Previously Described Subgrouping Rule: A Randomized Clinical Trial.
- Kankaanpaa M, Taimela S, Airaksinen O, et al. The efficacy of active rehabilitation in chronic low back pain. Effect on pain intensity, self-experienced disability, and lumbar fatigability. *Spine*. 1999;24(10):1034-42.
- Keller A, Brox JI, Gunderson R, et al. Trunk muscle strength, cross sectional area, and density in patients with chronic low back pain randomized to lumbar fusion or cognitive intervention and exercises. *Spine*. 2004;29(1):3-8.
- Kool J, et al. Exercise reduces sick leave in patients with non-acute non-specific low back pain: a meta-analysis. *J Rehabil Med*. 2004;36:49-62.

- Kroeling P, Gross A, Goldsmith CH, Burnie SJ, Haines T, Graham N, Brant A.  
Electrotherapy for neck pain. Cochrane Database of Systematic Reviews 2009, Issue 4.
- Kroeling P, Gross A, Graham N, et al. Electrotherapy for neck pain. Cochrane Database Syst Rev. 2013;(8):CD004251.
- Kumar S, Beaton K, Hughes T. The effectiveness of massage therapy for the treatment of nonspecific low back pain: A systematic review of systematic reviews. Int J Gen Med 2004;84(4):336-343.
- Kurtais Gursel Y, Ulus Y, Bilgic A, et al. Adding ultrasound in the management of soft tissue disorders of the shoulder: a randomized placebo-controlled trial. Phys Ther. 2004;84(4):336-343.
- Liebenson, C. The Chiropractic Rehabilitation Specialist and Quality Care. DC August 15, 1995, Volume 13, Issue 17.
- Little P, Lewith G, Webley F, et al. Randomised controlled trial of Alexander technique, exercise, massage (ATEAM) for chronic and recurrent back pain. BMJ 2008;337:a884.
- Lucy Beumer, Jennie Wong, Stuart J Warden, Joanne L Kemp, Paul Foster, Kay M Crossley. Effects of exercise and manual therapy on pain associated with hip osteoarthritis: a systematic review and meta-analysis. Br J Sports Med doi:10.1136/bjsports-2015-095255
- Macario A, Pergolizzi J. Systematic review of spinal decompression via motorized traction for chronic discogenic low back pain. Pain Practice. 2006;6(3):171-178.
- Manca A, Epstein DM, Torgerson DJ, Klaber Moffett JA, Coulton S, Farrin AJ, et al. Randomized trial of a brief physiotherapy intervention compared with usual physiotherapy for neck pain patients: cost-effectiveness analysis. Int J Technol Assess Health Care 2006;22:67-75.
- Maria Cristina Sandoval, Carolina Ramirez, Diana M Camargo, Tania F Salvini Effect of high-voltage pulsed current plus conventional treatment on acute ankle sprain.
- Mariénke van Middelkoop, Sidney M. Rubinstein, Ton Kuijpers, Arianne P. Verhagen, Raymond Ostelo, Bart W. Koes, and Maurits W. van Tulder. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. Eur Spine J. 2011 Jan; 20(1): 19–39.
- Miller J, Gross A, D'Sylva J, Burnie SJ, Goldsmith CH, Graham N, Haines T, Brønfort G, Hoving JL. Manual therapy and exercise for neck pain: a systematic review. Man Ther. 2010 Aug;15(4):334-54.
- Negrini S, Minozzi S, Bettany-Saltikov J, Zaina F, Chockalingam N, Grivas TB, Kotwicki T, Maruyama T, Romano M, Vasiliadis ES. Braces for idiopathic scoliosis in adolescents. Cochrane Database of Systematic Reviews 2010, Issue 1.

New Zealand Acute Low Back Pain Guide (June 2003 Edition) [www.nzgg.org.nz](http://www.nzgg.org.nz)

Niemisto L, Lahtinen-Suopanki T, Rissanen P, et al. A randomized trial of combined manipulation, stabilizing exercises, and physician consultation compared to physician consultation alone for chronic low back pain. *Spine*. 2003;28(19):2185-2191.

Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev*. 2014;10:CD011324.

Page MJ, Green S, Mrocki MA, et al. Electrotherapy modalities for rotator cuff disease. *Cochrane Database Syst Rev*. 2016;(6):CD012225.

Pia Damgaard, Else Marie Bartels, Inge Ris, Robin Christensen, and Birgit Juul-Kristensen. Evidence of Physiotherapy Interventions for Patients with Chronic Neck Pain: A Systematic Review of Randomised Controlled Trials. *ISRN Pain Volume 2013 (2013)*, Article ID 567175, 23 pages. <http://dx.doi.org/10.1155/2013/567175>

Poitras S, Brosseau L. Evidence-informed management of chronic low back pain with transcutaneous electrical nerve stimulation, interferential current, electrical muscle stimulation, ultrasound, and thermotherapy. *Spine Journal* 2008; 8(1):226-233.

Robertson VJ, Baker KG. A review of therapeutic ultrasound: effectiveness studies. *Phys Ther*. 2001 Jul;81(7):1339-50.

Roy Remmen, MD, PhD, Giannoula Tsakitzidis, PT, Wim Dankaerts, PT,MT, PhD, Paul Van Royen, MD, PhD. Non-Specific Neck Pain and Evidence-Based Practice. *European Scientific Journal* January 2013 edition vol.9, No.3 ISSN: 1857 – 7881 (Print) e - ISSN 1857- 7431

Rutjes AWS, Nüesch E, Sterchi R, Kalichman L, Hendriks E, Osiri M, Brosseau L, Reichenbach S, Jüni P. Transcutaneous electrostimulation for osteoarthritis of the knee. *Cochrane Database of Systematic Reviews* 2009, Issue 4.

Southerst, Danielle; Yu, Hainan; Randhawa, Kristi; Cote, Pierre; DAngelo, Kevin; et al. The effectiveness of manual therapy for the management of musculoskeletal disorders of the upper and lower extremities: a systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMA) Collaboration. *Chiropractic & Manual Therapies* 23:30 (2015).

Spitzer WO, LeBlanc FE, Dupuis M et al. Scientific approach to the assessment and management of activity-related spinal disorders. A monograph for clinicians. *Spine*. 1987;12:7S.

Susan Armijo-Olivo, Laurent Pitance, Vandana Singh, Francisco Neto, Norman, Thie, Ambra Michelotti. Effectiveness of Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders: Systematic Review and Meta-Analysis

Swenson RS. Therapeutic modalities in the management of nonspecific neck pain. *Phys Med Rehabil Clin N Am*. 2003; 14(3):605-27.

Syed Rehan Iftikhar Bukhari, Syed Shakil-ur-Rehman, Shakeel Ahmad, and Aamer Naeem. Comparison between effectiveness of Mechanical and Manual Traction combined with mobilization and exercise therapy in Patients with Cervical Radiculopathy *Pak J Med Sci*. 2016 Jan-Feb; 32(1): 31–34. doi: 10.12669/pjms.321.8923

Timm KE. A randomized-control study of active and passive treatments for chronic low back pain following L5 laminectomy. *J Orthop Sports Phys Ther*. 1994;20(6):276-286.

University of Michigan Health System. Acute low back pain guidelines for clinical care. [www.cme.med.umich.edu](http://www.cme.med.umich.edu). 2003.

van den Dolder PA, Ferreira PH, Refshauge KM. Effectiveness of soft tissue massage and exercise for the treatment of non-specific shoulder pain: A systematic review with meta-analysis. *Br J Sports Med* 2014;48(16):1216

van der Heijden G, Beurskens A, Koes B et al. The efficacy of traction for back and neck pain: a systematic blinded review of randomized clinical trial methods. *Phys Ther*. 1995;75:93–104.

van der Windt DAWM, van der Heijden GJ, Van den Berg S, ter Riet G, De Winter AF, Bouter LM, van den Bekerom MPJ. Therapeutic ultrasound for acute ankle sprains. *Cochrane Database of Systematic Reviews* 2002, Issue 1.

Walsh DM, Howe TE, Johnson MI, Sluka KA. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database of Systematic Reviews* 2009, Issue 2.

Wegner I, Widyahening IS, et al. Traction for low back pain with or without sciatica. *Cochrane Database Syst Rev*. 2013 Aug 19;8:CD003010. doi: 10.1002/14651858.CD003010.pub5.

Ylinen J, Kautiainen H, Wiren K, Hakkinen A. Stretching exercises vs manual therapy in treatment of chronic neck pain: a randomized, controlled cross-over trial. *J Rehabil Med*. 2007;39(2):126–132. Updated July 6, 2015

Zeng C, Li H, Yang T, et al. Electrical stimulation for pain relief in knee osteoarthritis: Systematic review and network meta-analysis. *Osteoarthritis Cartilage*. 2015;23(2):189-202.

Reviewed/Approved by  Michael Pentecost, MD, Chief Medical Officer