



Magellan Healthcare	
Clinical guidelines DURABLE MEDICAL EQUIPMENT	Original Date: August 2,2011 Page 1 of 9
	Adopted Date¹: April 4, 2016
Physical Medicine – Clinical Decision Making	Last Review Date: June 2017
Guideline Number: NIA_CG_609	Last Revised Date: June 2017
Responsible Department: Clinical Operations	Implementation Date: January 2018

Policy Statement

This policy will be used to define Durable Medical Equipment as well as support the medical necessity of the billed DME.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

Purpose

To outline the medical necessity of Durable Medical Equipment (DME).

Scope

This policy will apply to all physical medicine participating network practitioners, including chiropractors, physical therapists, occupational therapists, and speech language pathologists.

Definition

- Durable Medical Equipment (DME) is any equipment that provides therapeutic benefits to a patient for certain conditions and/or illnesses defined below
- DME consist of items which:
 - Are used to treat a defined illness or injury
 - Are not useful to a person in the absence of illness or injury
 - Are reusable and durable enough for repeated use
 - Are appropriate for use outside of a medical setting such as home, at school, or a work
- DME includes but is not limited to: back supports/braces, cervical collars, foot orthotics, electrical stimulation units, traction devices, and wheelchairs and assistive devices for gait.
- The use of any DME must have evidence of efficacy in the peer reviewed guideline, systematic review, and/or randomized controlled trial medical literature. The use of these devices is not considered medically necessary in the absence of scientific evidence in this peer reviewed medical literature.

¹ The “Original Date” above reflects the date the Policy was initiated by HSM Physical Health, Inc., (HSM). The “Adoption Date” above indicates the date that the Magellan Healthcare NIA Clinical Guideline Task Force reviewed and approved the Policy. HSM was acquired by National Imaging Associates, Inc., (NIA) in 2015 and is now a wholly owned subsidiary of NIA. National Imaging Associates, Inc., is a subsidiary of Magellan Healthcare, Inc.

Medical Necessity

Durable Medical Equipment and services are medically necessary when the following criteria are met:

1. The equipment is expected to provide improvement in specific, measurable functional deficits related to a documented illness or injury; **and**
2. The DME is provided by a health care professional; **and**
3. The equipment does not have significant non-medical uses; **and**
4. The clinical records clearly establish the medical need for the DME

Clinical documentation must include the following elements:

1. A diagnosis that justifies the equipment or supply being requested
2. A treatment plan (anticipated start and end date) for the use of the DME
3. Documented measurable functional deficit(s)
4. Expected outcomes and benefit related to a measurable functional deficit
5. Documentation of the healthcare providers training/education, supervision, and monitoring of the use of the DME as evidenced by the identification of provider type and signature in the record.
6. Documentation of a trial of conservative services that failed to improve a measurable functional deficit unless contraindicated
7. When appropriate, documentation of a trial of in-office care, such as cervical traction, that provided improvement in a measurable functional deficit
8. If an insurance plan does not cover a DME, then any visits associated with instruction on the DME would not be covered

DME

DME may be subject to medical necessity review. This would include: TENS or other electrical stimulation units, traction devices or chairs, etc. Additionally, any DME with a purchase or rental price of more than **\$200** will be subject to review.

Specific Durable Medical Equipment:

Electrical Stimulation for Pain

Transcutaneous electrical nerve stimulation (TENS) uses electrical stimulation at a painful site via the application of electrodes from the device to the surface of the skin. TENS devices generate electrical output, usually by a portable battery operated method.

Magellan considers TENS medically necessary DME when used as an adjunct or as an alternative to common conservative treatments for the initial 30 days of acute post-operative pain and for some forms of chronic musculoskeletal and neuropathic pain causing significant documented disruption of function unresponsive to at least a 1 month trial of conservative care including but not limited to manual therapy, active care, and pharmacotherapy. Please note that not all health plans reimburse for rental or purchase of home TENS units.

TENS is considered experimental and investigational for acute non-operative pain, acute and chronic headaches, deep abdominal pain, and chronic temporomandibular joint (TMJ) pain, adhesive capsulitis (frozen shoulder), chronic low back pain, neuropathic pain, pelvic

pain, phantom pain, stump pain, and all other indications because there is inadequate scientific evidence to support its efficacy for these specific types of pain.

A trial of TENS use for at least 30 days but not to exceed 90 days must be monitored by the healthcare provider. This trial period must include documentation of the effect on the patient's pain and measurable function to determine the effectiveness of the TENS unit. Treatment for long-term use is considered medically necessary if the trial period produced significant improvement in the patient's pain and measurable functional deficit(s). This documentation must include how the patient used the unit, the duration of use each time the unit was used, as well as the results of use. Concurrent chiropractic and/or physical therapy services are not indicated for the treatment of the same condition during the trial period.

Magellan does not consider the use of form-fitting conductive garments medically necessary.

Magellan considers the following forms of electrical stimulation not medically necessary: This list is not all-inclusive.

- Noninvasive neuron blockage devices
- Electroceutical therapy devices
- Bioelectric treatment systems
- Electro-Acuscope Therapy System
- Electrical stimulation of the sacral nerve roots or lumbosacral plexus for treatment of chronic pelvic or abdominal pain
- High-frequency pulsed electromagnetic stimulation
- Vagus nerve stimulation
- Bone growth stimulators

On June 8, 2012, the Centers for Medicare & Medicaid Services (CMS) rendered a decision memo for TENS for chronic low back pain. It states that TENS is not reasonable and necessary for the treatment of chronic low back pain.

In an evidence-based review, the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology evaluated the effectiveness of TENS in the treatment of pain in neurological disorders (Dubinsky and Miyasaki, 2010). There are conflicting reports of TENS compared to sham TENS in the treatment of chronic low back pain (LBP), with 2 Class II studies showing benefit, while 2 Class I studies and another Class II study not showing benefit. Because the Class I studies are stronger evidence, TENS is established as ineffective for the treatment of chronic LBP. The authors concluded that TENS is not recommended for the treatment of chronic LBP.

Guidelines on treatment of LBP from the National Collaborating Centre for Primary Care (Savigny et al, 2009) found insufficient evidence for the use of TENS in LBP and recommended against its use for that indication.

Evidence is insufficient to support the use of knee braces as a treatment for patellofemoral pain syndrome.

Home Traction Devices:

Home traction therapy is unproven and not medically necessary for treating low back and neck disorders with or without radiculopathy. The majority of studies are office based with mixed results. The quality of peer reviewed studies for home traction are limited as well to conclude that it is effective in the management of neck or low back pain or that it improves health outcomes. The indications for clinical application, patient selection criteria, risks, and comparison to alternative technologies have not been established for home traction therapy.

There is insufficient evidence from peer-reviewed published studies to conclude that lumbar spinal traction devices are effective at improving specific, measurable and functional deficits related to low back pain and leg-related low back pain. Magellan considers lumbar auto-traction devices experimental and investigational. This would include, but is not limited to: the Spinalator, the Arthrotonic stabilizer, the Anatomotor, Saunders Lumbar Hometrac, etc. Magellan also considers axial spinal uploading (gravity-dependent traction) devices experimental and investigation for the treatment of low back pain and leg-related low back pain. This would include, but is not limited to: the LTX 3000, VAX-D, and other decompression or traction devices, tables, weights or vests.

Orthotics, Prosthetics, Bracing and Assistive Devices

No definitive evidence as yet supports the use of orthoses in painful conditions of the cervical or lumbar spine. They should, therefore, be used only after individual consideration of the indications in each case.

Studies suggest that wearing a wrist splint can provide relief from carpal tunnel symptoms within a few weeks; however, the effect is only temporary.

The use of these devices must be necessary for the treatment of an illness or injury and to improve documented, measurable, and functional deficit(s). The documentation must include the reason the equipment is needed and the duration of its need.

A brace, orthotic, or prosthetic is a rigid or semi-rigid device. It is used to support and/or substitute a documented weak or deformed body part that is causing a documented measurable functional deficit.

The use of assistive devices is considered a standard of practice for general mobility needs and reduction in patients at risk of falling. Clinical documentation must support the use of these devices.

There is insufficient evidence to support the use of insoles or foot orthoses as either a treatment for LBP or in the prevention of LBP. Foot orthoses produce small short-term benefits in function and may also produce small reductions in pain for people with plantar fasciitis, but they do not have long-term beneficial effects compared with a sham device. Foot orthotics have no proven value for knee pain (other than medial osteoarthritis), pes planus (flat feet), pronation, corns and calluses, hip osteoarthritis, and lower leg injuries. Customized and prefabricated orthoses have similar effectiveness in the treatment of plantar fasciitis. Spinal Pelvic Stabilizers (Foot Levelers, Inc.) are specialized custom molded inserts designed to prevent foot injuries and improve foot alignment; these are considered experimental and investigational because their value in treatment of foot

disease has not been proven. The available evidence does not reveal any clear advantage of foot orthoses over simple insoles or physiotherapy for patellofemoral pain. While foot orthoses may help relieve knee pain over the short term, the benefit may be marginal. There is moderate evidence to support the use of foot orthotics in the treatment of chronic ankle instability to help improve postural control.

Overall, the evidence appears to suggest that custom foot orthotics and prefabricated orthotics have similar effectiveness. Therefore, prefabricated orthotics should be prescribed when there is a clinical indication for foot orthotics.

HCPCS 2017 Code L0631: Lumbar-sacral orthosis, sagittal control, with rigid anterior and posterior panels, posterior extends from sacrococcygeal junction to t-9 vertebra, produces intracavitary pressure to reduce load on the intervertebral discs, includes straps, closures, may include padding, shoulder straps, pendulous abdomen design, prefabricated item that has been trimmed, bent, molded, assembled, or otherwise customized to fit a specific patient by an individual with expertise. The clinical record must clearly document that this service involved customization of the lumbar orthosis in order for it to be reimbursable.

Strapping

The application of casts, splints, or straps is performed in an attempt to provide temporary immobilization or fixation to correct, protect, or stabilize a fracture, dislocation, or documented joint instability as a result of injury, disease or surgery.

The use of casting, splinting or strapping may be considered medically necessary for a patient who has experienced a fracture, dislocation or who has ligamentous instability following an acute injury, or as the result of a disease or surgery.

The application of casting, splinting or strapping **should not** be reported when any kind of treatment or restorative service aimed at correcting, protecting, or stabilizing the fracture, dislocation, or instability is concurrently done.

Strapping uses rigid material in order to restrict joint and/or muscular movement. Tape is typically worn for a relatively short duration (<18 hours). In contrast, kinesiology (kinesio) taping (KT) is a therapeutic taping method that utilizes a latex-free elastic tape, which is purported to give support and stability to joints and muscles without affecting circulation, range of motion, and biomechanics. It is also used for preventive maintenance, edema, and to treat pain. KT methods use highly-specific designed tape, which may be pre-cut for certain joints, and reportedly can be used by patients of every age and condition for 1-5 days per application. The consensus findings of peer reviews did not support the application of KT in clinical settings.

REFERENCES

- American College of Obstetricians and Gynecologists (ACOG). Chronic pelvic pain. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 2004 Mar. 17
- American College of Occupational and Environmental Medicine (ACOEM). Elbow disorders. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2007. 67 p
- American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology* 2010 Apr;112(4):810-33.
- Bigos SJ, Holland J, Holland C, Webster JS, Battie M, Malmgren JA. High-quality controlled trials on preventing episodes of back problems: systematic literature review in working-age adults. *Spine J.* 2009 Feb;9(2):147-68.
- Brosseau L, Wells GA, Finestone HM, Egan M, Dubouloz CJ, Graham I, Casimiro L, Robinson VA, Bilodeau M, McGowan J. Clinical practice guidelines for transcutaneous electrical nerve stimulation (TENS). *Top Stroke Rehabil* 2006 Spring;13(2):61-3.
- Cai C, Ming G, Ng LY. Development of a clinical prediction rule to identify patients with neck pain who are likely to benefit from home-based mechanical cervical traction. *Eur Spine J.* 2011;20(6):912-922
- Cancer pain management (general). In: Bader P, Echtler D, Fonteyne V, Livadas K, De Meerleer G, Paez Borda A, Papaioannou EG, Vranken JH. Guidelines on pain management. Arnhem, The Netherlands: European Association of Urology (EAU); 2010 Apr. p. 13-42.
- Carpal tunnel syndrome: Wrist splints and hand exercises. Institute for Quality and Efficiency in Health Care; Nov 5, 2014
- Centers for Medicare & Medicaid Services. Decision memo for transcutaneous electrical nerve stimulation for chronic low back pain (CAG-00429N). June 8, 2012. CMS: Baltimore, MD. Available at: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=256>.
- Chiropractic management of myofascial trigger points and myofascial pain syndrome: summary of clinical practice recommendations from the commission of the Council on Chiropractic Guidelines and Practice Parameters. Lexington, SC: Council on Chiropractic Guidelines and Practice Parameters (CCGPP); 2009.

- Chiropractic management of tendinopathy: summary of clinical practice recommendations from the commission of the Council on Chiropractic Guidelines and Practice Parameters. Lexington (SC): Council on Chiropractic Guidelines and Practice Parameters (CCGPP); 2009.
- Clarke JA, van Tulder MW, Blomberg SE, et al. Traction for low back pain with or without sciatica. *Cochrane Database Syst Rev.* 2007 Apr 18;(2):CD003010. Dallolio V. Lumbar spinal decompression with a pneumatic orthosis: Preliminary study. *Acta Neurochir Suppl.* Jan 1, 2005;92:133-7.
- Cruccu G, Aziz TZ, Garcia-Larrea L, Hansson P, Jensen TS, Lefaucheur JP, Simpson BA, Taylor RS. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol* 2007 Sep;14(9):952-70.
- Department of Veterans Affairs, Department of Defense. VA/DoD clinical practice guideline for rehabilitation of lower amputation. Washington (DC): Department of Veterans Affairs, Department of Defense; 2007. 163 p.
- Dubinsky RM, Miyasaki J. Assessment: Efficacy of transcutaneous electric nerve stimulation in the treatment of pain in neurologic disorders (an evidence-based review). Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology.* 2010;74(1):173-176.
- Gabriner ML, Braun BA, Houston MN, Hoch MC. The effectiveness of foot orthotics in improving postural control in individuals with chronic ankle instability: a critically appraised topic. *J Sport Rehabil.* 2015 Feb;24(1):68-71.
- Fall M., Baranowski AP., Elneil S., Engeler D., Hughes J., Messelink EJ., Oberpenning F., de C Williams AC., European Association of Urology (2010). EAU guidelines on chronic pelvic pain. [Eur Urol.](#) 2010 Jan;57(1):35-48. doi: 10.1016/j.eururo.2009.08.020. Epub 2009 Aug 31.
- Graham N, Gross A, Goldsmith CH, et al. Mechanical traction for neck pain with or without radiculopathy. *Cochrane Database Syst Rev.* 2008 Jul 16;(3):CD006408.
- Health Care Association of New Jersey (HCANJ). Pain management guideline. Hamilton (NJ): Health Care Association of New Jersey (HCANJ); 2006 Jul 18.
- Herr K, Bjoro K, Steffensmeier J, Rakel B. Acute pain management in older adults. Iowa City (IA): University of Iowa Gerontological Nursing Interventions Research Center, Research Translation and Dissemination Core; 2006 Jul.
- Hossain M, Alexander P, Burls A, Jobanputra P. Foot orthoses for patellofemoral pain in adults. *Cochrane Database Syst Rev.* 2011 Jan 19;(1):CD008402.
- Institute for Clinical Systems Improvement (ICSI). Assessment and management of chronic pain. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2009 Nov. 91 p.

Institute for Clinical Systems Improvement (ICSI). Diagnosis and treatment of headache. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2009 Mar. 76 p.

International Chiropractors Association of California. Management of whiplash associated disorders. Sacramento (CA): International Chiropractors Association of California; 2009. 55 p.

Khadilkar A, Odebiyi DO, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev.* 2008;(4):CD003008.

Kourosh Zarghooni, Frank Beyer, Jan Siewe, and Peer Eysel. The Orthotic Treatment of Acute and Chronic Disease of the Cervical and Lumbar Spine. *Dtsch Arztebl Int.* 2013 Nov; 110(44): 737–742.

Landorf KB, Keenan AM, Herbert RD. Effectiveness of Foot Orthoses to Treat Plantar Fasciitis: A Randomized Trial. *Arch Intern Med.* 2006;166(12):1305-1310.

Low back disorders. Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 2nd ed. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2007. 366 p.

Michigan Quality Improvement Consortium. Management of acute low back pain. Southfield (MI): Michigan Quality Improvement Consortium; 2010 Mar.

National Collaborating Centre for Chronic Conditions. Osteoarthritis. The care and management of osteoarthritis in adults. London (UK): National Institute for Health and Clinical Excellence (NICE); 2008 Feb. 22 p.

National Collaborating Centre for Primary Care. Low back pain. Early management of persistent non-specific low back pain. London (UK): National Institute for Health and Clinical Excellence (NICE); 2009 May. 25 p.

Nnoaham KE, Kumbang J. Transcutaneous electrical nerve stimulation (TENS) for 35 chronic pain. *Cochrane Database Syst Rev.* 2008:CD003222.

North American Spine Society (NASS). Diagnosis and Treatment of Cervical Radiculopathy from Degenerative Disorders; Burr Ridge (IL): North American Spine Society (NASS); 2010

North American Spine Society (NASS). Diagnosis and treatment of degenerative lumbar spinal stenosis. Burr Ridge (IL): North American Spine Society (NASS); Jan.2011

Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev.* 2014;10:CD011324.

Page MJ, Massy-Westropp N, O'Connor D, Pitt V. Splinting for carpal tunnel syndrome. *Cochrane Database Syst Rev.* 2012 Jul 11;(7):CD010003.

Parreira PdCS, Costa LdCM, Hespanhol Junior LC, Lopes AD, Costa LOP. Current evidence does not support the use of Kinesio Taping in clinical practice: a systematic review. *Journal of Physiotherapy* 60: 31–39.

Savigny P, Kuntze S, Watson P, et al. Low back pain: Early management of persistent non-specific low back pain. Full Guideline. London, UK: National Collaborating Centre for Primary Care and Royal College of General Practitioners; May 2009.

Sundstrøm T, Asbjørnsen H, Habiba S, Sunde GA, Wester K. Prehospital use of cervical collars in trauma patients: a critical review. *J Neurotrauma*. 2014 Mar 15;31(6):531-40.

Toward Optimized Practice. Guideline for the evidence-informed primary care management of low back pain. Edmonton (AB): Toward Optimized Practice; 2009 Mar 2. 21 p

University of Michigan Health System. Acute low back pain. Ann Arbor (MI): University of Michigan Health System; 2010 Jan. 14 p.

Vivienne Chuter, Martin Spink, Angela Searle, and Alan Ho. The effectiveness of shoe insoles for the prevention and treatment of low back pain: a systematic review and meta-analysis of randomised controlled trials. *BMC Musculoskelet Disord*. 2014; 15: 140.

Wegner I, Widyahening IS, van Tulder MW, Blomberg SE, de Vet HC, Brønfort G, et al. Traction for low-back pain with or without sciatica. *Cochrane Database Syst Rev*. 2013 Aug 19;8:

Zarghooni K, Beyer F, Siewe J, Eysel P. The orthotic treatment of acute and chronic disease of the cervical and lumbar spine. *Dtsch Arztebl Int*. 2013 Nov 1;110(44):737-42.

Reviewed/Approved by  Michael Pentecost, MD, Chief Medical Officer