



Magellan Healthcare	
Clinical guidelines ENDOMETRIAL CANCER	Original Date: June 2013 Page 1 of 6
Radiation Oncology	Last Review Date: July 2017
Guideline Number: NIA_CG_129	Last Revised Date: July 2017
Responsible Department: Clinical Operations	Implementation Date: January 2018

INTRODUCTION:

The majority of endometrial cancers are adenocarcinomas, with uterine sarcomas accounting for <10%. This clinical guideline will focus primarily on adenocarcinoma of the endometrium.

After a diagnosis of endometrial cancer is made, it is followed by a staging evaluation to determine extent of disease (local, regional, or metastatic), and prognostic findings. For patients in whom cancers of the uterus are suspected, an endometrial biopsy is typically performed. A review of the pathology will determine whether or not the tumors are of epithelial origin (endometrioid, papillary serous, clear cell, or carcinosarcoma) or stromal/mesenchymal carcinoma (stromal sarcoma or leiomyosarcoma). The majority of endometrial cancers, however, are adenocarcinomas with tumor typically confined to the uterus. Thus, this disease is often localized with an excellent prognosis. Current workup, including a complete surgical assessment, includes a histological grade, depth of myometrial invasion, and extent of extrauterine involvement. Prognostic factors are based on a pathologic assessment and include the percent of myometrial invasion, myometrial thickness, tumor size and location (upper fundus or lower uterine cervical), cervix involvement, and lymphovascular space involvement. The majority of patients are treated surgically with radiation reserved for patients who are deemed at a high risk of recurrence or for those deemed medically inoperable.

This guideline outlines several methods suitable for the employment of radiation therapy. This includes the use of 3-dimensional conformal radiation therapy and/or internal radiation (brachytherapy). IMRT is not indicated as a standard treatment option for uterine cancer. External beam treatments are typically delivered using a high-energy linear accelerator. Brachytherapy is generally delivered using temporary HDR sources such as iridium 192. The purpose of this guideline is to outline the most efficient, comparatively effective, diagnostic and treatment pathway. Treatment is typically broken down into patients in whom disease is limited to the uterus, cervical involvement (either suspected or confirmed), or extrauterine disease.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

INDICATIONS FOR RADIATION THERAPY AND TREATMENT OPTIONS:

Post-operative

- Brachytherapy Only (HDR or LDR, 5 fx maximum)
 - Stage IA – with adverse risk factors
 - Stage IA – without risk factors (Grades G2, 3)
 - Stage IB
 - Stage II – (Grade G1)
- External Beam Radiation Therapy Only (2D, 3D-CRT, 45-50.4 Gy, 28 fx maximum)
 - Stage IA – with adverse risk factors (Grades G2, 3)
 - Stage IB – without adverse risk factors (Grade G3)
 - Stage IB – with risk factors
 - Stage II – (Grade G1)
 - Stage III
 - Stage IV
- External Beam (2D, 3D-CRT, 45-50 Gy, 28 fx maximum) and Brachytherapy (HDR or LDR, 5 fx maximum)
 - Stage IA – with adverse risk factors (Grades G2, 3)
 - Stage IB – without risk factors (Grade G3)
 - Stage IB – with risk factors
 - Stage II – (Grades G1, 2, 3)
 - Stage IIIA & IIIB & IIIC (Grades G1, 2, 3)

Medically Inoperable/ Pre-Operative

- Brachytherapy Only (HDR or LDR, 7 fx maximum)
 - Stage I & II
- External Beam Radiation Therapy Only (2D, 3D-CRT, 45-50 Gy, 28 fx maximum)
 - All Stages
- External Beam (2D, 3D-CRT, 45-50.4 Gy) and Brachytherapy (HDR or LDR, 4 fx maximum)
 - All Stages

Palliative

- Up to 10 fx

Unless otherwise indicated standard radiation fractionation consists of 1.8 Gy to 2.0 Gy per day.

TREATMENT OPTIONS REQUIRING ADDITIONAL CLINICAL REVIEW:

Intensity Modulated Radiation Therapy (IMRT)

IMRT is not indicated as a standard treatment option and should not be used routinely for the delivery of radiation therapy for endometrial cancer. IMRT is strictly defined by the utilization of inverse planning modulation techniques. IMRT may be appropriate for limited circumstances in which radiation therapy is indicated and 3D conformal radiation therapy (3D-CRT) techniques cannot adequately deliver the radiation prescription without exceeding normal tissue radiation tolerance, the delivery is anticipated to contribute to potential late toxicity or tumor volume dose heterogeneity is such that unacceptable hot or cold spots are created.

Clinical rationale and documentation for performing IMRT rather than 2D or 3D-CRT treatment planning and delivery will need to:

- Demonstrate how 3D-CRT isodose planning cannot produce a satisfactory treatment plan (as stated above) via the use of a patient specific dose volume histograms and isodose plans.
- Provide tissue constraints for both the target and affected critical structures.

Stereotactic Body Radiation Therapy (SBRT)

Stereotactic Body Radiation Therapy is not a standard treatment option for the treatment of endometrial cancer.

Proton Beam Radiation Therapy

Proton beam is not an approved treatment option for endometrial cancer. Proton beam has not been proven superior treatment to conventional radiation therapy.

REFERENCES

- Alektiar, K.M., Venkatraman, E., Chi, D.S., & Barakat, R.R. (2005). Intravaginal brachytherapy alone for intermediate risk endometrial cancer. *Int J Radiat Oncol Biol Phys.* 62(1), 111-117. doi:10.1016/j.ijrobp.2004.09.054.
- American College of Radiology. ACR Appropriateness Criteria (2014). Advanced Stage Endometrial Cancer. Access online at: <https://acsearch.acr.org/docs/3083084/Narrative/> on June 12, 2015.
- ASTEC/EN.5 Study Group, Blake, P., Swart, A.M., et al. (2009). Adjuvant external beam radiotherapy in the treatment of endometrial cancer (MRS ASTEC and NCIC CTG EN.5 randomized trials): pooled trial results, systematic review, and meta-analysis. *Lancet.* 373(9658), 137-146. doi: 10.1016/S0140-6736(08)61767-5.
- American Society for Therapeutic Radiology and Oncology (ASTRO). Cervical and endometrial cancer patients report fewer side effects and better quality of life with IMRT. September 26, 2016. Accessed on May 15, 2017 at https://www.astro.org/uploadedFiles/MAIN_SITE/News_and_Publications/News/News_Releases/2016/ASTRO_2016_Klopp.pdf
- Bentzen SM, Constine LS, Deasy JO, et al. Quantitative analyses of normal tissue effects in the clinic QUANTEC: An introduction to the scientific issues. Introductory paper. *Int J Radiat Oncol Biol Phys.* 2010; 76(3):S3-S9.
- Choosing Wisely. Keyword: Endometrial. Accessed on May 15, 2017 at [http://www.choosingwisely.org/clinician-lists/#keyword=Endometrial&parentSociety=American Society for Radiation Oncology](http://www.choosingwisely.org/clinician-lists/#keyword=Endometrial&parentSociety=American_Society_for_Radiation_Oncology)
- Creutzberg, C.L., van Putten, W.L., Koper, P.C., et al. (2000). Surgery and Postoperative radiotherapy versus surgery alone for patients with stage I endometrial carcinoma: multicentre randomized trial. PORTEC Study Group. *Post Operative Radiation Therapy in Endometrial Carcinoma.* 355, 1404-1411. Retrieved from [http://dx.doi.org/10.1016/S0140-6736\(00\)02139-5](http://dx.doi.org/10.1016/S0140-6736(00)02139-5).
- Creutzberg, C.L., van Putten, W.L., Warlam-Rodenhuis, C.C., et al. (2004). Outcome of high-risk stage IC, grade 3, compared with stage I endometrial carcinoma patients: the Postoperative Radiation Therapy Endometrial Carcinoma Trial. *J Clin Oncol.* 22, 1234-1241. doi: 10.1200/JCO.2004.08.159.
- Homesley, H.D., Filiaci, V., Gibbons, S.K., et al. (2008). Randomized phase III trial in advanced endometrial carcinoma of surgery and volume-directed radiation followed by cisplatin and doxorubicin with or without paclitaxel: A Gynecologic Oncology Group study [abstract]. 39th Annual Meeting of the Society of Gynecologic Oncologists, Tampa, FL, Plenary Session I, 9 March 2008 Abstract #1. *Gynecol Oncol.* 108, S2. Retrieved from [http://www.google.com/search?q=Willett,+C.G.,+%26+Gunderson,+L.L.+\(2007\).+Perez+and+Brady%E2%80%99s+principles+and+practice+of+radiation+oncology,+5th+ed.+Lipp](http://www.google.com/search?q=Willett,+C.G.,+%26+Gunderson,+L.L.+(2007).+Perez+and+Brady%E2%80%99s+principles+and+practice+of+radiation+oncology,+5th+ed.+Lipp)

[incott+Williams+%26+Wilkins.+1318-1335.&safe=active&rls=com.microsoft:en-us:IE-Address&rlz=1I7ADBF&ei=6TKkUdu6LpLk8gTKs4CIDA&start=10&sa=N](#)

Jackson A, Marks LB, Bentzen SM, et al. The lessons of QUANTEC: Recommendations for reporting and gathering data on dose-volume dependencies of treatment outcome. *Int J Radiat Oncol Biol Phys.* 2010; 76(3):S155-S160.

Johnson, N., & Comes, P. (2007). Survival and recurrent disease after Postoperative radiotherapy for early endometrial cancer: Systematic review and meta-analysis. *BJOG* 114(11), 1313-1320. DOI: 10.1111/j.1471-0528.2007.01332.x.

Keys, H.M., Roberts, J.A., Brunetto, V.L., et al. (2004). A phase III trial of surgery with or without adjunctive external pelvic radiation therapy in intermediate risk endometrial adenocarcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol.* 92, 744-751. Retrieved from <http://dx.doi.org/10.1016/j.ygyno.2003.11.048>.

Klopp, Ann, et. al. (2014) The Role of Postoperative Radiation Therapy for Endometrial Cancer: An ASTRO Evidence-Based Guideline. *Practical Radiation Oncology*. Accessed online at: <http://www.practicalradonc.org/cms/attachment/2014676400/2036188629/mmc1.pdf> on June 2, 2015.

Koh, W.J., Tran, A.B., Douglas, J.G., et al. (2001). Radiation therapy in endometrial cancer. *Best Pract Res Clin Obstet Gynaecol.* 15, 417-432. doi:10.1053/beog.2001.0186.

Kong, A., Johnson, N., Comes, P., et al. (2007). Adjuvant radiotherapy for stage I endometrial cancer. *Chochrane Database Syst Rev.* 2, CD003916.

Kong, A., Powell, M., Blake, P. (2008). The role of Postoperative radiotherapy in carcinoma of the endometrium. *Clin Oncol (R Coll Radiol).* 20(6), 457-462. Retrieved from <http://dx.doi.org/10.1016/j.clon.2008.03.011>.

Lee, C.M., Szabo, A., Shrieve, D.C., et al. (2006). Frequency and effect of adjuvant radiation therapy among women with stage I endometrial adenocarcinoma. *JAMA.* 295, 389-397. doi:10.1001/jama.295.4.389.

National Comprehensive Cancer Network (NCCN). Uterine Neoplasms Version 2.2017. Retrieved May 15, 2017 from: https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf

Nout, R.A., Putter, H., Jurgenliemk-Schulz, I.M., et al. (2008). Vaginal brachytherapy versus external beam pelvic radiotherapy for high-intermediate risk endometrial cancer. Results of the randomized PORTEC-2 trial [abstract]. *J Clin Oncol.* 26, LBA5503. doi: 10.1016/S0140-6736(09)62163-2.

Portelance, L. et al: A Phase II Multi-institutional Study Of Postoperative Pelvic Intensity Modulated Radiation Therapy (IMRT) With Weekly Cisplatin In Patients With Cervical Carcinoma: Two Year Efficacy Results Of The RTOG 0418 Proceedings of the American

Society for Radiation Oncology 54th Annual Meeting. Vol 84, No. 3 Nov 1 2012. *Int J Radiat Oncol Biol Phys*.

Randall, M.E., Filiaci, V.L., Muss, H., et al. (2006). Randomized phase III trial of whole-abdominal irradiation versus doxorubicin and cisplatin chemotherapy in advanced endometrial carcinoma: A Gynecologic Oncology Group Study. *J Clin Oncol*. 24, 36-44. doi: 10.1200/JCO.2004.00.7617.


Roeske JC, et al: Intensity-modulated whole pelvic radiation therapy in patients with gynecologic malignancies. *Int J Radiat Oncol Biol Phys*. 2000 Dec 1;48(5):1613-21.

Scholten, A.N., van Putten, W.L., Beerman, H., et al. (2005). PORTEC Study Group Postoperative radiotherapy for stage I endometrial carcinoma: long-term outcome of the randomized PORTEC trial with central pathology review. *Int J Radiat Oncol Biol Phys*. 63(3), 834-838. doi:10.1016/j.ijrobp.2005.03.007.

Shih KK et al: Postoperative pelvic intensity-modulated radiotherapy in high risk endometrial cancer. *Gynecol Oncol*. 2013 Mar;128(3):535-9. doi: 10.1016/j.ygyno.2012.11.020. Epub 2012 Nov 20.

Solhjem, M.C., Petersen, I.A., & Haddock, M.G. (2005). Vaginal brachytherapy alone is sufficient adjuvant treatment of surgical stage I endometrial carcinoma. *Int J Radiat Oncol Biol Phys*. 62(5), 1379-1384. doi:10.1016/j.ijrobp.2005.01.026.

Susumu, N., Sagae, S., Udagawa, Y., et al. (2008). Randomized phase III trial of pelvic radiotherapy versus cisplatin-based combined chemotherapy in patients with intermediate and high risk endometrial cancer: A Japanese Gynecologic Oncology Group study. *Gynecol Oncol*. 108(1), 226-233. Retrieved from <http://dx.doi.org/10.1016/j.ygyno.2007.09.029>.

Reviewed/Approved by  Michael Pentecost, MD, Chief Medical Officer