



Magellan Healthcare	
Clinical guidelines: NON-CANCEROUS CONDITIONS	Original Date: March 2015 Page 1 of 6
Radiation Oncology	Last Review Date: July 2017
Guideline Number: NIA_CG_135	Last Revised Date: July 2017
Responsible Department: Clinical Operations	Implementation Date : January 2018

INTRODUCTION:

Radiation therapy may have appropriate use in several non-malignant conditions. The treatment goal in patients with non-malignant conditions is to achieve relief of the indicated condition with radiation therapy with minimal risk of radiation exposure to sensitive structures.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

INDICATIONS FOR RADIATION THERAPY

2 D or 3D Conformal (3D CRT) is considered medically necessary for several non-malignant conditions including but not limited to:

- Prevention of keloid scars as an adjunctive therapy following excisional surgery
- Heterotopic ossification
- Pterygium in cases that cannot be medically managed
- Villonodular synovitis

Stereotactic Radiation Therapy (SRS, SBRT) is considered medically necessary when used in the treatment of non-malignant cranial lesions including the following:

- Arteriovenous malformation (AVM) of the brain or spine.
- Trigeminal neuralgia that has not responded to other, more conservative, treatments.
- Non cancerous brain tumors such as acoustic neuroma, benign schwannomas, meningioma, hemangioma, pituitary adenoma, craniopharyngioma, neoplasm of the pineal gland, and chordomas

Also refer to NIA Stereotactic Radiation Therapy Guideline.


TREATMENT OPTIONS REQUIRING PHYSICIAN REVIEW:

Treatment for other non-malignant conditions utilizing proton beam, stereotactic radiation therapy (SBRT), or intensity modulated radiation therapy (IMRT) modalities should be referred to physician review.

REFERENCES:

- Aqqarwal, A., Fersht, N., Brada, M. (2013) Radiotherapy for craniopharyngioma. Pituitary. Mar;16(1):26-33. doi: 10.1007/s11102-012-0429-1.
- Bentzen SM, Constine LS, Deasy JO, et al. Quantitative analyses of normal tissue effects in the clinic QUANTEC: An introduction to the scientific issues. Introductory paper. Int J Radiat Oncol Biol Phys. 2010; 76(3):S3-S9.
- Casentini, L., Fornezza, U., Perini, Z., Perissinotto, E., Columbo, F. (2015) Multisession stereotactic radiosurgery for large vestibular schwannomas. J Neurosurg. 16:1-7.
- Combs, S. E., Engelhand, C., Kopp, C., Wiedenmann, N., Schramm, O., Prokic, V., ...Grosu, A. L. (2015). Long-term outcome after highly advanced single-dose or fractionated radiotherapy in patients with vestibular schwannomas - Pooled results from 3 large German centers. Radiother Oncol. Mar;114(3):378-83. doi: 10.1016/j.radonc.2015.01.011.
- Ding, D., Yen, C. P., Starke, R. M., Lee, C. C., Sheehan, J. P. (2014) Unyielding progress: recent advances in the treatment of central nervous system neoplasms with radiosurgery and radiation therapy. J Neurooncol. 119(3):513-29. doi: 10.1007/s11060-014-1501-7.
- Flickinger, J. C. (2011) A radiobiological analysis of multicenter data for postoperative keloid radiotherapy. Int J Radiat Oncol Biol Phys. Mar 15;79(4):1164-70. doi: 10.1016/j.ijrobp.2009.12.019.
- Gross, C. E., Frank, R. M., Hsu, A. R., Diaz, A. Gitelis, S. (2015) External Beam Radiation Therapy for Orthopaedic Pathology. J Am Acad Orthop Surg. 2015 Apr;23(4):243-252
- Hasan, S., Young, M., Albert, T., Shah, A. H., Okoye, C., Bregy, A., ...Komotar, R. J. (2015). The Role of Adjuvant Radiotherapy After Gross Total Resection of Atypical Meningiomas. World Neurosurg. Dec 19. pii: S1878-8750(14)01408-9. doi: 10.1016/j.wneu.2014.12.037.
- Jackson A, Marks LB, Bentzen SM, et al. The lessons of QUANTEC: Recommendations for reporting and gathering data on dose-volume dependencies of treatment outcome. Int J Radiat Oncol Biol Phys. 2010; 76(3):S155-S160.
- Kondziolka, D., Perez, B., Flickinger, J. C., Habeck, M., Lunsford, L. D., (1998). Arch Neurol. 55(12):1524-1529. Gamma knife radiosurgery for trigeminal neuralgia: Results and expectations.
- Maesawa, S., Salame, C., Flickinger, J. C., Pirris, S., Kondziolka, D., Lunsford, L. D. (2001). Clinical outcomes after stereotactic radiosurgery for idiopathic trigeminal neuralgia. J Neurosurg. 94(1):14-20

- Maniakas, A., Saliba, I. (2012) Microsurgery versus stereotactic radiation for small vestibular schwannomas: a meta-analysis of patients with more than 5 years' follow-up. *Otol Neurotol.* 33(9):1611-1620.
- Paryani, S. B., Scott, W. P., Wells, J.W. Jr., Johnson, D. W., Chobe, R. J., Kuruvilla, A., Schoepel, S., Deshmukh, A. (1994) Management of pterygium with surgery and radiation therapy. The North Florida Pterygium Study Group. *Int J Radiat Oncol Biol Phys.* Jan 1;28(1):101-3.
- Pashtan, I., Oh, K. S., Loeffler, J. S. (2014) Radiation therapy in the management of pituitary adenomas. *Handb Clin Neurol.* 124:317-24. doi: 10.1016/B978-0-444-59602-4.00021-6
- Popovic, M., Aqarwal, A., Zhang, L, Yip, C., Kreder, H. J., Nousiainen, M. T., Jenkinson, R., ...Chow, E. (2014). Radiotherapy for the prophylaxis of heterotopic ossification: a systematic review and meta-analysis of published data. *Radiother Oncol.* 2014 Oct;113(1):10-7. doi: 10.1016/j.radonc.2014.08.025.
- Portnow, L. H., Scott, M. , Morris, C. G., Mendenhall, W. M., Marcus, R. B., Indelicato, D. J. (2012) Fractionated radiotherapy in the management of benign vascular tumors. *Am J Clin Oncol.* Dec;35(6):557-61. doi: 10.1097/COC.0b013e31821f847f
- Seregard, S., Pelayes, D. E., Singh, A. D. (2013) Radiation therapy: uveal tumors. *Dev Ophthalmol.* 52:36-57. doi: 10.1159/000351055.
- Seegenschmiedt, M. H., Micke, O., Muecke, R., & the German Cooperative Group on Radiotherapy for Non-malignant Diseases (GCG-BD). (2015). Radiotherapy for non-malignant disorders: state of the art and update of the evidence-based practice guidelines. *The British Journal of Radiology*, 88(1051), 20150080. <http://doi.org/10.1259/bjr.20150080>. Accessed on May 15, 2017 from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4628533/>
- Sonier, M., Gete, E., Herbert, C., McKenzie, M., Murphy, J., Moiseenko, V. (2014). Intensity-modulated stereotactic radiosurgery for arteriovenous malformations: guidance for treatment planning. *Radiat Oncol.* Mar 10;9:73. doi: 10.1186/1748-717X-9-73.

Reviewed/Approved by  Michael Pentecost, MD, Chief Medical Officer