



<b>National Imaging Associates, Inc.</b>	
<b>Clinical guidelines PLAIN FILM X-RAYS</b>	<b>Original Date:        Adopted 2016</b>
<b>Physical Medicine – Clinical Decision Making</b>	<b>Last Revised Date:    July 2019</b>
<b>Guideline Number: NIA_CG_610</b>	<b>Implementation Date: January 2020</b>

### **Policy Statement**

The use of plain films is medically necessary when clinical findings dictate their utilization. Films are not indicated to identify unsuspected contraindications to chiropractic manipulation, view postural changes and biomechanics, or identify subluxations. Insufficient scientific evidence exists to support the use of routine plain film radiographs as a means for improved clinical outcomes in spinal disorders. There is insufficient clinical research to support improved clinical outcomes when radiographs are a part of a routine component of the initial evaluation or ongoing treatment. This organization has adopted the Diagnostic Imaging Practice Guidelines for Musculoskeletal Complaints in Adults. These guidelines represent the official position of the Council on Chiropractic Guidelines and Practice Parameters in matters relating to the use of diagnostic imaging in the chiropractic profession.

The use of full spine radiographs, except for the clinical investigation and diagnosis of scoliosis, is not supported by clinical research.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

### **Purpose**

This policy will be used to support the medical necessity of plain film radiographs by chiropractic providers within the first 30 days of care.

### **Scope**

This policy will apply to all participating network chiropractic practitioners.

### **Definition**

#### Plain films:

Spinal or extremity radiographs used as a diagnostic tool by chiropractors.

### **Guidelines:**

- An appropriate history and examination are required to identify if plain films are clinically indicated.

- Utilization of radiographs by chiropractors will not be reimbursed unless sufficient medical record documentation is submitted with claims to support the medical necessity of the film. The clinical record must clearly document the rationale for the x-rays, any suspected pathology, or what condition the chiropractor hopes to rule out. The use of plain films to rule out an unsuspected pathology is not clinically indicated.
- Routine use of radiographs as part of the initial evaluation or part of an ongoing treatment plan will not be reimbursed.
- The use of full spine radiographs for any diagnosis other than scoliosis is not considered medically necessary and will not be reimbursed.
- Contraindications to plain film x-rays include:
  - Infants (0-36 months)
  - Pregnancy or possible pregnancy
  - Obesity, if size precludes good radiographic resolution
  - Patient has positioning difficulty due to mental status or physical restrictions, which precludes good radiographic resolution
  - Children 3 to 18 years of age, except for investigation of suspected acute fracture, dislocation, infection, scoliosis, developmental defects, or a suspected pathology.
- Requirements:
- The clinical record must contain a written x-ray report within five (5) business days from the date of service.
- The clinic must have all of the following:
  - A documented Quality Control Program inclusive of both imaging equipment and film processors
  - A documented Radiation Safety and As Low As Reasonably Achievable (ALARA) Program
  - Documented emergency policies, procedures, and equipment on site (i.e. automated external defibrillator (AED))
  - Documentation of current Basic Life Support (BLS) certification
  - Records of formal preventative maintenance program per original equipment specifications
  - A current (within three (3) years) letter of state inspection, calibration report, or physicist's report
  - At a minimum, an automatic processor must be used to develop all analog plain films.

#### **CLINICAL EXAMPLES of MEDICALLY NECESSARY X-RAYS:**

- Investigation of suspected acute fracture
- Follow up radiographs to monitor a healing fracture
- Investigation of suspected bony dislocation
- Evaluation of prior surgical site where manual based treatment may be applied (where no previous films are available for review)

- Suspect (patient history, pain characteristics and/or physical examination) malignancy, infection, systemic disease, or inflammatory spondyloarthropathy
- Precise quantification of clinically suspected active child or juvenile scoliosis
- Persistent (same or worse pain) after first month of treatment
- Significant history of drug or alcohol abuse such as IV drugs or chronic alcoholism or chronic use of steroids
- Adult patient with thoracolumbar, lumbar, or thoracic spine blunt trauma or acute injuries (falls, motor vehicle accidents [MVA's], motorcycle, pedestrian, cyclists, etc)
- Adults with complicated (ie, "red flag") LBP, thoracic pain, or neck pain & indicators of contraindication to SMT (relative/absolute)
- Suspected inflammatory spondyloarthropathies, neoplasia, or infection
- Adult patient: in the absence of expected treatment response or worsening after 4 weeks of conservative treatment
- Adult patient with acute neck injury and positive CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients.)
- Suspected lumbar degenerative spinal stenosis or spondylolisthesis if patient is greater than 50 years of age and/or has progressive neurological deficit – AP (or PA) and lateral lumbar views
- Adult with recent unimaged blunt trauma to pelvis and unable to bear weight – AP pelvis and lateral hip "frog leg" views
- Acute neck pain with recent unimaged dangerous trauma, paresthesia in extremities or age greater than 65 or non-traumatic neck pain with radicular symptoms – APOM, AP lower cervical and lateral neutral views
- Adult with painful or progressive scoliosis – Erect standing full spine (14x36) PA and lateral views in the absence of recent films. Plain film x-rays may be appropriate when red flags suggest further screening for cancer, infection, or fracture. They may also be sufficient for the initial evaluation of patients with the following red flags: age >70 years, a history of recent significant trauma, or risk of osteoporosis. Plain film x-rays may be appropriate, but are usually not sufficient for clinical decision making without advanced imaging, in the presence of other red flags. Radiographs are unreliable for assessment of bone mass changes before at least a 30%-50% loss. In healthy peri- and early postmenopausal women (45-64 yoa), consider using the OST score (Osteoporosis Self Assessment Tool.) OST score considers only 2 variables: (weight in kg – age)/5. The cut-off for a positive test is <2, indicating this woman should be referred for DXA.
- Current x-ray recommendations/guidelines for spinal and extremity disorders emphasize a focused history and physical examination, reassurance, initial pain management medications if necessary (acetaminophen or nonsteroidal anti-inflammatory drugs), and consideration of nonpharmacologic therapies (e.g., manipulation, exercise, etc.) without routine imaging in patients with nonspecific neck and/or low back pain (Bussi eres, 2007, 2008; Dagenais, 2010; Koes, 2010; Pillastrini, 2011). Imaging is considered for those without improvement after 6 weeks and for those with clinical indicators of serious pathologies (red flags) (Bach, 2009; Bussi eres, 2007, 2008; Chou, 2011, 2007).

Plain film x-rays of the extremities may be indicated in the following circumstances:

- Significant history of recent trauma sufficient to cause fracture
- Significant history of repetitive stress to cause stress fracture
- History or clinical findings of malignancy
- Previous surgery or fracture
- Suspicion of or confirmed inflammatory arthritis

- Evaluation of gross deformities
- Bruising, swelling, redness heat, indicating infection
- Lymphadenopathy
- Evaluation of developmental hip dysplasia in the pediatric population
- Evaluation of Leg-Calve-Perthes disease
- Evaluation of slipped capital femoral epiphysis in the pediatric population

Plain film radiographs may be appropriate but are usually not sufficient for clinical decision making without advanced imaging (MR and/or CT) in the presence of other red flags including [Davis, Dagenais (2012)]:

- Age < 20 years or > 50 years
- Failure to improve with care, no prior films
- Personal history of intravenous drug abuse
- History of malignancy
- Immune suppression
- Night pain
- Night pain (unrelated to movement)
- Pain at multiple sites
- Pain at rest
- Fever
- Structural deformity
- Systemic unwellness
- Unexplained weight loss

Spinal radiographs also have a role in evaluation of scoliosis and in postoperative evaluation of instrumentation and fusion (Davis, 2011). For the evaluation of scoliosis in children, radiographic decision-making and examinations should be performed in accordance with guidance published by the American College of Radiology (ACR) and the Society for Pediatric Radiology (SPR) (Faerber, 2009, 2012). Radiographic examination is indicated for pediatric patients at high risk for cervical spine instability – especially those with Down syndrome (Faerber, 2012).

**INITIAL PLAIN FILM X-RAYS ARE NOT INDICATED in the FOLLOWING CASES:**

- Adult patient with acute uncomplicated LBP (<4 wks' duration). Uncomplicated definition: nontraumatic pain without neurologic deficits or indicators of potentially serious pathologies
- Adult patient with uncomplicated subacute (4-12 wks' duration) or persistent LBP (>12 wks' duration) AND no previous treatment trial
- Adult patient with nontraumatic acute LBP (<4 wks' duration) AND sciatica and no red flags
- Sciatica, unless patient is age >50 or has progressive neurological deficits
- Suspected lumbar disc herniation
- Suspected degenerative spondylolithesis/lateral stenosis, unless patient is age >50 or has progressive neurological deficits
- Suspected lumbar degenerative spinal stenosis, unless patient is age >50 or has progressive neurological deficits
- Adult patient with uncomplicated acute (<4 wk duration) thoracic spine pain

- Adult patient with uncomplicated subacute (4-12 wk duration) or persistent (>12 week duration) thoracic spine pain and no previous treatment trial
- Adult patient with nonpainful and nonprogressive scoliosis
- Adult patient with acute uncomplicated neck pain (<4 wks' duration)
- Adult patient with uncomplicated subacute neck pain (4-12 wks' duration) with or without arm pain
- Adult patient with persistent neck pain (>12 wk) with or without arm pain
- Adult patient with acute neck injury and negative CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients) (Stiell 2011)  
<http://jama.jamanetwork.com/article.aspx?articleid=194296>.
- In headache complaints, vital signs (to R/O severe hypertension or fever) and testing of the cranial nerves (to R/O vascular events, space occupying lesions, etc) should be an integral part of initial examination. Significant positive findings mandate further evaluation. Without red flags or significant findings, no initial films are indicated.
- Coccyx trauma and coccydynia
- The routine use of spinal radiographs for structural and biomechanical analysis has not been substantiated to improve patient outcomes (Peterson 2005). The clinical evidence is insufficient to support an association between sagittal (lordosis, kyphosis) spinal curves and health outcomes including spine-related pain (Christensen 2008). The utility of plain film radiography for the detection of spinal 'subluxations', or to guide the specifics of spinal manipulative therapy, is controversial (Petersen 2005). "The validity of the various systems of roentgenometric analysis has not been proven and their underlying premise of bilateral symmetry within the body does not take into account natural structural anomalies" (Petersen 2005). Adding to this controversy is the fact that nonspecific spinal abnormalities are common in asymptomatic patients (Davis 2011).
- "Strong evidence shows that routine back imaging does not improve patient outcomes, exposes patients to unnecessary harms, and increases costs" (Chou 2012). "Available evidence indicates that immediate, routine lumbar spine imaging in patients with LBP and without features indicating a serious underlying condition, did not improve outcomes compared with usual clinical care without immediate imaging. Clinical care without immediate imaging seems to result in no increased odds of failure in identifying serious underlying conditions in patients without risk factors for these conditions. In addition to lacking clinical benefit, routine lumbar imaging is associated with radiation exposure (radiography and CT) and increased direct expenses for patients and may lead to unnecessary procedures. This evidence confirms that clinicians should refrain from routine, immediate lumbar imaging in primary care patients with nonspecific, acute or subacute LBP and no indications of underlying serious conditions" (Andersen 2011).

**POLICY HISTORY:**

**Review Date:** July 23, 2019

**Review Summary:**

- Format updated for improved readability

## REFERENCES

American College of Radiology (ACR). *ACR–SPR practice guideline for the performance of radiography for scoliosis in children*. ACR. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Scoliosis.pdf>. Revised 2014. Retrieved April 9, 2018.

American College of Radiology (ACR). *ACR–ASSR–SPR–SSR practice guideline for the performance of spine radiography*. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Rad-Spine.pdf>. Revised 2017. Retrieved April 9, 2018.

Ammendolia C, Hogg-Johnson S, Pennick V, et al. Implementing evidence-based guidelines for radiography in acute low back pain: A pilot study in a chiropractic community. *J Manip Physiol Ther*. 2004; 27(3):170-179.

Andersen JC. Is immediate imaging important in managing low back pain? *J Athl Train*. 2011; 46:99-102.

Bach SM, Holten KB. Guideline update: What's the best approach to acute low back pain? *J Fam Pract*. 2009; 58:E1.

Bigos S, Bowyer O, Brean G, et al. *Clinical practice guideline No. 14.: Acute low back problems in adults*. Washington, DC: US Government Printing Office; AHCPR Publication No. 95-0642; 1994.

Borenstein DG. A clinician's approach to acute low back pain. *Am J Med*. 1997; 102(1A):16S-22S.

Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging guideline for musculoskeletal complaints in adults – an evidence-based approach: Introduction. *J of Manip Physiol Ther*. 2007a; 30:617-683.

Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging guideline for musculoskeletal complaints in adults – an evidence-based approach – part 2: upper extremity disorders. *J Manip Physiol Ther*. 2008a; 31:2-32.

Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging guideline for musculoskeletal complaints in adults – an evidence-based approach – part 3: spinal disorders. *J Manip Physiol Ther*. 2008b; 31:31-88.

Bussièrès AE, Taylor JA, Peterson C. Diagnostic imaging guideline for musculoskeletal complaints in adults – an evidence-based approach – part 1: lower extremity disorders. *J Manip Physiol Ther*. November-December 2007a; 30(9):684-717.

Chou R, Deyo RA, Jarvik JG. Appropriate use of lumbar imaging for evaluation of low back pain. *Radiol Clin N Am*. 2012; 50:569-585.

Chou R, Qaseem A, Owens DK, Shekelle P. Diagnostic imaging for low back pain: Advice for high-value health care from the American College of Physicians. *Ann Intern Med*. 2011; 154:181-189.

Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: A joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007; 147:478-491.

Christensen ST, Hartvigsen J. Spinal curves and health: A systematic critical review of the epidemiological literature dealing with associations between sagittal spinal curves and health. *J Manip Physiol Therap.* 2008; 31(9):690-714.

Dagenais S, Tricco AC, Haldeman S. Assessment of Low Back Pain. In: Dagenais S, Haldeman S, eds. *Evidence-Based Management of Low Back Pain.* St Louis, Missouri. Mosby (Elsevier) 2012:21-31.

Dagenais S, Tricco AC, Haldeman S. Synthesis of recommendations for the assessment and management of low back pain from recent clinical practice guidelines. *Spine J.* 2010; 10:514-529.

Davis PC, Wippold II FJ, Cornelius RS, et al. American College of Radiology ACR Appropriateness Criteria<sup>®</sup> - Low Back Pain. <http://www.acr.org/~media/ACR/Documents/AppCriteria/Diagnostic/LowBackPain.pdf>. Published 2011.

Faeber EN, Disler DG, Hernanz-Schulman M, et al. American College of Radiology ACR–SPR practice guideline for the performance of radiography for scoliosis in children. <http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/Scoliosis.pdf>. Revised 2009.

Faeber EN, Kan JH, Newman B, et al. ACR–ASSR–SPR–SSR practice guideline for the performance of spine radiography. American College of Radiology. [http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/Spine\\_Radiography.pdf](http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/Spine_Radiography.pdf). Revised 2012.

Henschke N, Maher CG, Ostelo RWJG, et al. Red flags to screen for malignancy in patients with low-back pain. *Cochrane Database Syst Rev.* 2013;(2): CD008686. doi: 10.1002/14651858.CD008686.pub2.

Institute for Clinical Systems Improvement (ICSI). *Healthcare guidelines, low back pain, adult.* September 2003. <https://www.icsi.org/>.

Jarvik JG. Imaging of adults with low back pain in the primary care setting. *Neuroimaging Clin N Am.* 2003; 13(2):293-305.

Kendrick D, Fielding K, Bentley E, et al. Radiography of the lumbar spine in primary care patients with low back pain: randomized controlled trial. *BMJ.* 2001; 322:400-405.

Kendrick D, Fielding K, Bentley E, et al. The role of radiography in primary care patients with low back pain of at least 6 weeks duration: A randomized controlled trial. *Health Technol Assess.* 2001; 5(30):1-69.

Koes BW, van Tulder M, Lin CW, et al. An updated overview of clinical guidelines for the management of non-specific low back pain in primary care. *Eur Spine J.* December 2010; 19(12): 2075-2094.

Magora A, Schwartz A. Relation between low back pain syndrome and X-ray findings. Degenerative osteoarthritis. *Scand J Rehabil Med.* 1976; 8:115-25.

Mootz R, Hoffman L, Hansen D. Optimizing clinical use of radiology and minimizing radiation exposure in chiropractic practice. *Topics in Clinical Chiropractic.* 1997; 4:34-44.



New Zealand Guidelines Group. *New Zealand Acute Low Back Pain Guide*. October 2004. <https://www.healthnavigator.org.nz/media/1006/nz-acute-low-back-pain-guide-acc.pdf>. Retrieved April 9, 2018.

Patel ND, Broderick DF, Burns J, et al. ACR Appropriateness Criteria Low Back Pain. *J Am Coll Radiol*. September 2016; 13(9):1069-1078.

Petersen C, Hsu W. Indications for and use of x-rays. In: Haldeman SC, ed. *Principles and Practice of Chiropractic*. 3<sup>rd</sup> ed. McGraw-Hill Education; 2005:661-681.

Pillastrini P, Gardenghi I, Bonetti F, et al. An updated overview of clinical guidelines for chronic low back pain management in primary care. *Joint Bone Spine*. 2011; <http://dx.doi.org/10.1016/j.jbspin.2011.03.019>.

Stiell I, Wells G, Vandemheen K, et al. The Canadian c-spine rule for radiography in alert and stable trauma patients. *JAMA*. 2001; 286(15):1841-1848.

Tran B, Saxe JM, Ekeh AP. Are flexion extension films necessary for cervical spine clearance in patients with neck pain after negative cervical CT scan? *J Surg Res*. September 2013; 184(1):411-413.

Van den Hoogen HMM, Koes BW, van Eijk JT, et al. On the accuracy of history, physical examination, and erythrocyte sedimentation rate in diagnosing low pain in general practice. A criteria-based review of the literature. *Spine*. 1995; 20:318-27.


Van Tulder MW, Tuut M, Pennick V, et al. Quality of primary care guidelines for acute low back pain. *Spine*. 2004; 29(17):E357-362.

Williams CM, Henschke N, Maher CG, et al. Red flags to screen for vertebral fracture in patients presenting with low-back pain. *Cochrane Database Syst Rev*. 2013; (1): CD008643. doi: 10.1002/14651858.CD008643.pub2.

Wipf JF, Deyo RA. Low back pain. *Med Clin North Am*. 1995; 79:231-246.

Yelland M. Diagnostic Imaging for Back Pain. *Aust Fam Physician*. June 2004; 33(6):415-419.

Young KJ. Historical influence on the practice of chiropractic radiology: part I - a survey of diplomates of the American Chiropractic College of Radiology. *Chiropr Man Therap*. 2017; 25:14.

Reviewed / Approved by  Patrick Browning, VP, Medical Director

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.