

<b>National Imaging Associates, Inc.*</b>	
<b>Clinical guidelines PANCREATIC CANCER</b>	<b>Original Date: June 2013</b>
<b>Radiation Oncology</b>	<b>Last Revised Date: February 2020</b>
<b>Guideline Number: NIA_CG_134</b>	<b>Implementation Date: January 2021</b>

#### INDICATIONS FOR RADIATION THERAPY:

2D and 3D conformal radiation therapy techniques are considered medically necessary for treatment of pancreatic cancer.

#### **Neoadjuvant (Pre- Operative) or Resectable or Borderline Resectable without evidence of metastatic (NCCN, 2019)**

- No standard treatment regimen currently exists for this subset of patients. If neoadjuvant radiation therapy is delivered, a dose of 45-54 Gy in 1.8-2.5 Gy fractions or 36 Gy in 2.4 fractions are viable options.

#### **Adjuvant (Post-Operative) Resectable Without Evidence of Metastatic Disease (NCCN, 2019)**

- For resected cases (45-46 Gy in 1.8-2 Gy fractions) to the clinical target volume, followed by boost (5-9Gy). Up to 31 fractions.

#### **Unresectable/Locally Advanced Without Evidence of Metastatic Disease (NCCN, 2019)**

- Radiation delivered in 45-54 Gy (1.8-2.5 Gy fractions or 36 Gy in 2.4 fractions). Up to 30 fractions.

#### **Palliative (NCCN, 2019)**

- Radiation delivered in 25-36 Gy in 2.4-5.0 Gy fractions is usual for patients with metastatic disease who require palliation for obstruction or pain. Up to 15 fractions.

#### **Local Recurrence after Resection without Evidence of Systemic Metastatic Disease**

- Adjuvant chemotherapy or chemoradiation if no previous radiation given. Up to 30 fractions. (NCCN, 2019)

#### TREATMENT OPTIONS REQUIRING PHYSICIAN REVIEW:

##### **Intensity Modulated Radiation Therapy (IMRT)**

IMRT is not indicated as a standard treatment option and should not be used routinely for the delivery of radiation therapy for pancreatic cancer. IMRT is strictly defined by the utilization of inverse planning modulation techniques. IMRT may be appropriate for limited circumstances in which radiation therapy is indicated and 3D conformal radiation therapy (3D-CRT) techniques cannot adequately deliver the radiation

\* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

prescription without exceeding normal tissue radiation tolerance, the delivery is anticipated to contribute to potential late toxicity or tumor volume dose heterogeneity is such that unacceptable hot or cold spots are created.

Clinical rationale and documentation for performing IMRT rather than 2D or 3D-CRT treatment planning and delivery will need to:

- Demonstrate how 3D-CRT isodose planning cannot produce a satisfactory treatment plan (as stated above) via the use of patient specific dose volume histograms and isodose plans.
- Provide tissue constraints for both the target and affected critical structures.

### **Stereotactic Body Radiation Therapy (SBRT) (NCCN, 2019)**

*Stereotactic Body Radiation Therapy (SBRT) is considered medically necessary for the treatment of Pancreatic Cancer. If requested, a physician review is required.*

### **Proton Beam Radiation Therapy**

Proton beam is not an approved treatment option for pancreatic cancer. Proton beam has not been proven a superior treatment to conventional radiation therapy.

### **Intra Operative Radiation Therapy (IORT)**

The role of interoperative radiation therapy for pancreatic cancer is controversial but may be reasonable for patients undergoing resection that may result in closer involved margins. IORT may be considered on a case by case basis.

---

### **BACKGROUND:**

Pancreatic cancer typically occurs later in life. Risk factors include smoking, alcohol use, obesity, diabetes, and certain chemical exposures. Pancreatitis has also been shown to have an increased risk of developing pancreatic cancer. Surgical resection is potentially the only curative approach, but most patients present with more advanced stage disease. Overall, the actuarial five-year survival rate is approximately 20%.

The goal of these guidelines is to delineate appropriate indications of the employment of radiation therapy in the treatment of pancreatic cancer and to define suitable methods of delivery of radiation therapy for these indications.

### **POLICY HISTORY:**

**Review Date:** February 2019

**Review Summary:** Added and updated references

**Review Date:** February 2020

**Review Summary:**

- Stereotactic Radiation Therapy:  
Deleted: Stereotactic Body Radiation Therapy (SBRT) is not currently an approved treatment option for pancreatic cancer.

Added: Stereotactic Body Radiation Therapy (SBRT) is considered medically necessary for the treatment  
Pancreatic Cancer

- Added and Updated References

## REFERENCES:

- American College of Radiology (ACR). ACR Appropriateness Criteria®. Borderline and unresectable pancreas cancer. <https://acsearch.acr.org/docs/3096669/Narrative/>. Published 2016. Accessed May 2, 2018.
- American College of Radiology (ACR). ACR Appropriateness Criteria®. Resectable pancreas cancer. <https://acsearch.acr.org/docs/3097847/Narrative/>. Published 2016. Accessed May 15, 2017.
- American Society for Therapeutic Radiology and Oncology (ASTRO). Chemotherapy and stereotactic ablative radiation (SABR) consecutively may be promising treatment option for patients with advanced pancreatic cancer. [https://www.astro.org/News-and-Publications/News-and-Media-Center/News-Releases/2014/Chemotherapy-and-stereotactic-ablative-radiation-\(SABR\)-consecutively-may-be-promising-treatment-option-for-patients-with-advanced-pancreatic-cancer/](https://www.astro.org/News-and-Publications/News-and-Media-Center/News-Releases/2014/Chemotherapy-and-stereotactic-ablative-radiation-(SABR)-consecutively-may-be-promising-treatment-option-for-patients-with-advanced-pancreatic-cancer/). Published September 15, 2014. Accessed May 15, 2017.
- Artinyan A, Anaya DA, McKenzie S, et al. Neoadjuvant therapy is associated with improved survival in resectable pancreatic adenocarcinoma. *Cancer*. 2011; 117:2044-2049. doi: 10.1002/cncr.25763.
- Bentzen SM, Constine LS, Deasy JO, et al. Quantitative analyses of normal tissue effects in the clinic QUANTEC: An introduction to the scientific issues. Introductory paper. *Int J Radiat Oncol Biol Phys*. 2010; 76(3):S3-S9.
- Chang DT, Schellenberg D, Shen J, et al. Stereotactic radiotherapy for unresectable adenocarcinoma of the pancreas. *Cancer*. February 1, 2009; 115(3):665-672. doi: 10.1002/cncr.24059.
- Corsini MM, Miller RC, Haddock MG, et al. Adjuvant radiotherapy and chemotherapy for pancreatic carcinoma: the Mayo Clinic experience (1975-2005). *J Clin Oncol*. 2008; 26:3511-3516. doi: 10.1200/JCO.2007.15.8782.
- Crane CH, Beddar AS, Evans DB. The role of intraoperative radiotherapy in pancreatic cancer. *Surg Oncol Clin N Am*. 2003; 12:965-977. [http://www.ncbi.nlm.nih.gov/pubmed/?term=Surg+Oncol+Clin+N+Am.+2003+Oct%3B12\(4\)%3A965-77](http://www.ncbi.nlm.nih.gov/pubmed/?term=Surg+Oncol+Clin+N+Am.+2003+Oct%3B12(4)%3A965-77).
- Evans DB, Varadhachary GR, Crane CH, et al. Preoperative gemcitabine-based chemoradiation for patients with resectable adenocarcinoma of the pancreatic head. *J Clin Oncol*. July 20, 2008; 26(21):3496-3502. doi: 10.1200/JCO.2007.15.8634.
- Feng M, Balter JM, Normolle D, et al. Characterization of pancreatic tumor motion using cine MRI: Surrogates for tumor position should be used with caution. *Int J Radiat Oncol Biol Phys*. July 1, 2009; 74(3):884-891. doi: 10.1016/j.ijrobp.2009.02.003.
- Ford EC, Herman J, Yorke E, et al. 18F-FDG PET/CT for image-guided and intensity-modulated radiotherapy. *J Nucl Med*. October 2009; 50(10):1655-1665. doi: 10.2967/jnumed.108.055780.
- Goldstein SD, Ford EC, Duhon M, et al. Use of respiratory-correlated four-dimensional computed tomography to determine acceptable treatment margins for locally advanced pancreatic adenocarcinoma. *Int J Radiat Oncol Biol Phys*. February 1, 2010; 76(2):597-602. doi: 10.1016/j.ijrobp.2009.06.009.

Heinrich S, Pestalozzi B, Lesurtel M, et al. Adjuvant gemcitabine versus Neoadjuvant gemcitabine/oxaliplatin plus adjuvant gemcitabine in resectable pancreatic cancer: A randomized multicenter phase III study (NEOPAC study). *BMC Cancer*. 2011; 11:346. doi: 10.1186/1471-2407-11-346.

Herman JM, Swartz MJ, Hsu CC, et al. Analysis of fluorouracil-based adjuvant chemotherapy and radiation after pancreaticoduodenectomy for ductal adenocarcinoma of the pancreas: Results of a large, prospectively collected database at the Johns Hopkins Hospital. *J Clin Oncol*. July 20, 2008; 26(21):3503-3510. doi: 10.1200/JCO.2007.15.8469.

Huguet F, Girard N, Guerche CS, et al. Chemoradiotherapy in the management of locally advanced pancreatic carcinoma: A qualitative systematic review. *J Clin Oncol*. May 1, 2009; 27(13):2269-2277. doi: 10.1200/JCO.2008.19.7921.

Jackson A, Marks LB, Bentzen SM, et al. The lessons of QUANTEC: Recommendations for reporting and gathering data on dose-volume dependencies of treatment outcome. *Int J Radiat Oncol Biol Phys*. 2010; 76(3):S155-S160.

Koong AC, Christofferson E, Le QT, et al. Phase II study to assess the efficacy of conventionally fractionated radiotherapy followed by a stereotactic radiosurgery boost in patients with locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys*. October 1, 2005; 63(2):320-323. [http://www.redjournal.org/article/S0360-3016\(05\)01153-3/abstract](http://www.redjournal.org/article/S0360-3016(05)01153-3/abstract).

Koong AC, Christofferson E, Le QT, et al. Phase I study of stereotactic radio-surgery in patients with locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys*. 2004; 58(4): 1017-1021. <http://med.stanford.edu/stanfordhospital/PDF/cyberknife/phase2PancreasIJROBP.pdf>.

Landry J, Catalano PJ, Staley C, et al. Randomized phase II study of gemcitabine plus radiotherapy versus gemcitabine, 5-fluorouracil, and cisplatin followed by radiotherapy and 5-fluorouracil for patients with locally advanced, potentially resectable pancreatic adenocarcinoma. *J Surg Oncol*. 2010; 101:587-592. doi: 10.1002/jso.21527.

Laurence JM, Tran PD, Morarji K. A systematic review and meta-analysis of survival and surgical outcomes following neoadjuvant chemoradiotherapy for pancreatic cancer. *J Gastrointest Surg*. 2011; 15:2059-2069. doi: 10.1007/s11605-011-1659-7.

Le Scodan R, Mornex F, Girard N, et al. Preoperative chemoradiation in potentially resectable pancreatic adenocarcinoma: Feasibility, treatment effect evaluation and prognostic factors, analysis of the SFRO-FFCD 9704 trial and literature review. *Ann Oncol*. August 2009; 20(8):1387-1396. doi: 10.1093/annonc/mdp015.

Marti JL, Hochster HS, Hiotis SP, et al. Phase I/II trial of induction chemotherapy followed by concurrent chemoradiotherapy and surgery for locoregionally advanced pancreatic cancer. *Ann Surg Oncol*. 2008; 15:3521-3531. doi: 10.1245/s10434-008-0152-3.

Massucco P, Capussotti L, Magnino A, et al. Pancreatic resections after chemoradiotherapy for locally advanced ductal adenocarcinoma: analysis of perioperative outcome and survival. *Ann Surg Oncol*. 2006; 13:1201-1208. <http://link.springer.com/article/10.1245%2Fs10434-006-9032-x>.

McClaine RJ, Lowy AM, Sussman JJ, et al. Neoadjuvant therapy may lead to successful surgical resection and improved survival in patients with borderline resectable pancreatic cancer. *HPB (Oxford)*. 2010; 12:73-79. doi: 10.1111/j.1477-2574.2009.00136.x.

Minn AY, Schellenberg D, Maxim P, et al. Pancreatic tumor motion on a single planning 4D-CT does not correlate with intrafraction tumor motion during treatment. *Am J Clin Oncol*. August 2009; 32(4):364-368. doi: 10.1097/COC.0b013e31818da9e0.

National Comprehensive Cancer Network (NCCN). Pancreatic Adenocarcinoma. Version 1.2019. 20[https://www.nccn.org/professionals/physician\\_gls/pdf/pancreatic.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf). Accessed February 12, 2020.

Neoptolemos JP, Stocken DD, Bassi C, et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs. gemcitabine following pancreatic cancer resection: A randomized controlled trial. *JAMA*. 2010; 304:1073-1081. doi: 10.1001/jama.2010.

Patel M, Hoffe S, Malafa M, et al. Neoadjuvant GTX chemotherapy and IMRT-based chemoradiation for borderline resectable pancreatic cancer. *J Surg Oncol*. August 1, 2011; 104(2):155-161. doi: 10.1002/jso.21954.

Pawlik TM, Laheru D, Hruban RH, et al. Evaluating the impact of a single-day multidisciplinary clinic on the management of pancreatic cancer. *Ann Surg Oncol*. August 2008; 15(8):2081-2088. doi: 10.1245/s10434-008-9929-7.

Reviewed / Approved by  Sri Gorty, M.D., Physician Clinical Reviewer, Radiation Oncology

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.