

Managed Health Services

Specific policy administered by National Imaging Associates, Inc. (NIA)

Clinical guidelines	
Limited or Localized Follow Up Sinus CT	
CPT Codes: 76380	Original Date: September 1997
Guideline Number: Managed Health Services_CG_009	Last Revised Date (by Managed Health Services): May 2023
	Last Reviewed Date (by NIA): May 2023
	Implementation Date: June 2023

INDICATIONS FOR SINUS & MAXILLOFACIAL CT

Rhinosinusitis¹⁻⁵

- Clinical suspicion of fungal infection^{6, 7}
- Clinical suspicion of complications,⁸ such as
 - Preseptal, orbital, or intracranial infection⁹
 - Osteomyelitis
 - Cavernous sinus thrombosis
- Acute (<4weeks) or subacute (4-12 weeks) sinusitis (viral or bacterial)
 - Not responding to medical management including 2 or more courses of antibiotics at least 5 days each course

Note: Imaging may be indicated in those predisposed to complications, including diabetes, immune-compromised state, or a history of facial trauma or surgery.

- Recurrent acute rhinosinusitis with 4 or more annual episodes without persistent symptoms in between and is a possible surgical candidate
- Chronic recurrent sinusitis (>12 weeks) not responding to medical management*, is a possible surgical candidate, and with at least two of the following:
 - mucopurulent discharge
 - nasal obstruction and congestion
 - facial pain, pressure, and fullness
 - decreased or absent sense of smell

*Note: Medical management for chronic sinusitis includes nasal saline irrigation and/or topical intranasal steroids. In chronic sinusitis, repeat imaging is not necessary unless clinical signs or symptoms have changed.

1— Managed Health Services specific policy administered by NIA

- Allergic Rhinitis – sinus imaging usually not indicated unless there are signs of complicated infection, signs of neoplasm, or persistence of symptoms/chronic rhinosinusitis despite treatment (including antihistamines) and is a possible surgical candidate¹⁰
- If suspected as a cause of poorly controlled asthma (endoscopic sinus surgery improves outcomes)¹¹
- To evaluate in the setting of unilateral nasal polyps or obstruction (to evaluate for a potential neoplasm)³

Pediatrics Rhinosinusitis^{12, 13}

- Persistent or recurrent sinusitis not responding to treatment (primarily antibiotics, treatment may require a change of antibiotics)
- Suspicion of orbital or central nervous system involvement (e.g., swollen eye, proptosis, altered consciousness, seizures, nerve deficit)
- Clinical suspicion of a fungal infection (more common in immunocompromised children)

Deviated nasal septum, polyp, or other structural abnormality seen on imaging or direct visualization

- Causing significant airway obstruction AND
- Imaging is needed to plan surgery or determine if surgery is appropriate^{14, 15}

Suspected sinonasal mass based on exam, nasal endoscopy, or prior imaging with contraindication to MRI or if bony involvement suspected^{3, 16, 17}

Refractory Asthma - these patients benefit from medical treatment and surgery together^{11, 18, 19}

Anosmia or Dysosmia noted on objective testing, is persistent, of unknown origin and MRI cannot be performed^{16, 17, 20, 21}

Suspected infection

- Osteomyelitis (after x-rays and MRI cannot be performed)²²
- Abscess based on clinical signs and symptoms of infection

Face mass^{16, 17, 23}

- Present on physical exam and remains non-diagnostic after x-ray or ultrasound is completed; **OR**
- Known or highly suspected head and neck cancer on examination¹⁶; **OR**
- Failed 2 weeks of treatment for suspected infectious adenopathy²⁴

Facial trauma²⁵⁻³⁰

- Severe facial trauma
- Suspected facial bone fracture with indeterminate x-ray
- For further evaluation of a known fracture for treatment or surgical planning

2— Managed Health Services specific policy administered by NIA

LIMITED OR LOCALIZED FOLLOW UP SINUS CT

- CSF (cerebrospinal fluid) rhinorrhea when looking to characterize a bony defect
Note: for CSF otorrhea should be a Temporal Bone CT; for intermittent leaks and complex cases, consider CT/MRI/Nuclear Cisternography. CSF fluid should always be confirmed with laboratory testing (Beta-2 transferrin assay)

Salivary gland

- Sialadenitis (infection and inflammation of the salivary glands) with indeterminate ultrasound, bilateral symptoms or concern for abscess³¹
- Suspected or known salivary gland stones³²⁻³⁴

Granulomatosis with polyangiitis (Wegener's granulomatosis) disease³⁵

Suspected Osteonecrosis of the Jaw³⁶

- Possible etiologies: bisphosphonate treatment, dental procedures, Denosumab, radiation treatment

Lesion seen on x-ray or other study requiring further characterization (primary or secondary bone tumor, metabolic disorder)³⁷

Trigeminal neuralgia/neuropathy if MRI is contraindicated or cannot be performed (for evaluation of the extracranial nerve course)

- If atypical features (i.e., bilateral, hearing loss, dizziness/vertigo, visual changes, sensory loss, numbness, pain > 2min, pain outside trigeminal nerve distribution, progression)^{6, 38}

Pre-operative/procedural evaluation

- Pre-operative evaluation for a planned surgery or procedure

Post-operative/procedural evaluation

- When imaging, physical, or laboratory findings indicate surgical or procedural complications

COMBINATION OF STUDIES WITH SINUS & MAXILLOFACIAL CT

Sinus CT/Chest CT

- Granulomatosis with polyangiitis (Wegener's granulomatosis) disease (GPA)³⁹

BACKGROUND

Computed tomography (CT) primarily provides information about bony structures but may also be useful in evaluating soft tissue masses. It can help document the extent of facial bone fractures, facial infections, and abscesses, and can aid in diagnosing salivary stones.

Additionally, CT may be useful in characterizing and identifying tumor extent in the face and may be used in the assessment of chronic osteomyelitis.

CT scans can provide more detailed information about the anatomy and abnormalities of the paranasal sinuses than plain films. A CT scan provides greater definition of the sinuses and is more sensitive than plain radiography for detecting sinus pathology, especially within the sphenoid and ethmoid sinuses. CT scan findings can be nonspecific, however, and should not be used routinely in the diagnosis of acute sinusitis. The primary role of CT scans is to aid in the diagnosis and management of recurrent and chronic sinusitis, or to define the anatomy of the sinuses prior to surgery.

CT vs MRI - MRI allows better differentiation of soft tissue structures within the sinuses. It is used occasionally in cases of suspected tumors or fungal sinusitis. Otherwise, MRI has no advantages over CT scanning in the evaluation of sinusitis. Disadvantages of MRI include high false-positive findings, poor bony imaging, and higher cost. MRI scans take considerably longer to accomplish than CT scans and may be difficult to obtain in patients who are claustrophobic.

Rhinosinusitis - Society consensus recommendation is not to order sinus computed tomography (CT) or indiscriminately prescribe antibiotics for uncomplicated acute rhinosinusitis.⁴² Viral infections cause the majority of acute rhinosinusitis and only 0.5 percent to 2 percent progress to bacterial infections. Most acute rhinosinusitis resolves without treatment in two weeks. Uncomplicated acute rhinosinusitis is generally diagnosed clinically and does not require a sinus CT scan or other imaging. Antibiotics are not recommended for patients with uncomplicated acute rhinosinusitis who have mild illness and assurance of follow-up. If a decision is made to treat, amoxicillin should be first-line antibiotic treatment for most acute rhinosinusitis.

Anosmia - Nonstructural causes of anosmia include post viral symptoms, medications (Amitriptyline, Enalapril, Nifedipine, Propranolol, Penicillamine, Sumatriptan, Cisplatin, Trifluoperazine, Propylthiouracil). These should be considered prior to advanced imaging to look for a structural cause. Anosmia and dysgeusia have been reported as common early symptoms in patients with COVID-19, occurring in greater than 80 percent of patients. For isolated anosmia, imaging is typically not needed once the diagnosis of COVID has been made, given the high association. As such, COVID testing should be done prior to imaging.^{20, 40, 41}

Suspected Osteonecrosis of the Jaw - CT can characterize the extension of the lesions and in detecting cortical involvement. MRI should be reserved for those patients who have soft tissue extension of the disease.⁴²

Trigeminal Neuralgia - According to the International Headache Society, TN is defined as “a disorder characterized by recurrent unilateral brief electric shock-like pain, abrupt in onset and termination, limited to the distribution of one or more divisions of the trigeminal nerve and triggered by innocuous stimuli.”⁴³

POLICY HISTORY

Date	Summary
March 2022	<p>Reformatted and update references Reformatted and updated background Reformatted-structural abnormality, salivary gland, and trauma sections Clarified:</p> <ul style="list-style-type: none"> • Sialadenitis (infection and inflammation of the salivary glands) with indeterminate ultrasound, bilateral symptoms, or concern for abscess • acute vs subacute sinusitis • described medical management for acute (including 2 or more courses of antibiotics at least 5 days each course) and chronic sinusitis (includes nasal saline irrigation and/or topical intranasal steroids) • Abscess <p>Added:</p> <ul style="list-style-type: none"> • Note: Imaging may be indicated in those predisposed to complications, including diabetes, immune-compromised state, or a history of facial trauma or surgery (Acute sinusitis) • And is a surgical candidate- for chronic sinusitis and recurrent acute rhinosinusitis • In chronic sinusitis, repeat imaging is not necessary unless clinical signs or symptoms have changed. • Indications for allergic rhinitis <p>Removed:</p> <ul style="list-style-type: none"> • 4 weeks of medical management for acute and chronic sinusitis
April 2021	<p>Updated background section and references Added:</p> <ul style="list-style-type: none"> • Chronic recurrent sinusitis (symptoms for >12 weeks) not responding to at least 4 weeks of medical management and with at least two of the following: <ul style="list-style-type: none"> ○ mucopurulent discharge ○ nasal obstruction and congestion ○ facial pain, pressure, and fullness ○ decreased or absent sense of smell • Facial Trauma- For further evaluation of a known fracture for treatment or surgical planning • Suspected sinonasal mass based on exam, nasal endoscopy, or prior imaging with contraindication to MRI or if bony involvement suspected • Dysosmia • Sialadenitis with indeterminate ultrasound or bilateral symptoms

	<p>Clarified:</p> <ul style="list-style-type: none"> • Rhinosinusitis - Symptoms that persist for more than 4 weeks and are not responding to medical management (e.g., 2 or more courses of antibiotics or any combination of antibiotics, steroids or antihistamines for more than 4 weeks) • CSF (cerebrospinal fluid) rhinorrhea <i>when looking to characterize a bony defect</i> (for CSF otorrhea should be a Temporal Bone CT; for intermittent leaks and complex cases consider CT/MR/Nuclear Cisternography). CSF fluid should always be confirmed with laboratory testing (Beta-2 transferrin assay) • Suspicion of salivary gland stones <p>Deleted:</p> <ul style="list-style-type: none"> • For poorly controlled asthma associated with upper respiratory tract infection. May be performed without failing 4 consecutive weeks of treatment with medication. • Trigeminal neuralgia – if Age < 40
<p>May 2020</p>	<ul style="list-style-type: none"> • Updated references; Updated and reordered background information • Reordered and reformatted indications • Clarified: <ul style="list-style-type: none"> ○ Reworded: Rhinosinusitis: Clinical suspicion of complications, such Preseptal, orbital or intracranial infection, Osteomyelitis, Cavernous sinus thrombosis ○ Deviated nasal septum, polyp, or other structural abnormality seen on imaging or direct visualization that may be causing significant airway obstruction (if needed to plan surgery or determine if surgery is appropriate) ○ Refractory Asthma (Sinus CT) - these patients benefit from medical treatment and surgery together ○ Anosmia noted on objective testing, is persistent, of unknown origin and MRI cannot be done ○ Suspected infection: Osteomyelitis (after x-rays, MRI cannot be done) <p style="margin-left: 40px;">Facial trauma: Post traumatic CSF (cerebrospinal fluid) rhinorrhea (for CSF otorrhea should be a Temporal Bone CT)</p> • Added: <ul style="list-style-type: none"> • Rhinosinusitis <ul style="list-style-type: none"> ○ Recurrent acute rhinosinusitis with 4 or more annual episodes without persistent symptoms in between ○ If suspected as a cause of poorly controlled asthma (endoscopic sinus surgery improves outcomes) (Vashishta, 2013)

	<ul style="list-style-type: none"> ○ To evaluate in the setting of unilateral nasal polyps or obstruction (to evaluate for a potential neoplasm) (Rosenfeld, 2015) ● Pediatrics Rhinosinusitis (ACR, 2018; Wald, 2013) <ul style="list-style-type: none"> ○ Persistent or recurrent sinusitis not responding to treatment (primarily antibiotics, treatment may require a change of antibiotics) ○ Suspicion of orbital or central nervous system involvement (e.g., swollen eye, proptosis, altered consciousness, seizures, nerve deficit) (Ward, 2013) ○ Clinical suspicion of a fungal infection (more common in immunocompromised children). <p>Added:</p> <ul style="list-style-type: none"> ● Suspected Osteonecrosis of the Jaw (Popovic, 2010) <ul style="list-style-type: none"> ○ Possible etiologies: bisphosphonate treatment, dental procedures, Denosumab, radiation treatment) ○ CT can characterize the extension of the lesions and in detecting cortical involvement. MRI should be reserved for those patients who have soft tissue extension of the disease ● Lesion seen on x-ray or other study requiring further characterization (primary or secondary bone tumor, metabolic disorder) ● Trigeminal neuralgia/neuropathy if MRI is contraindicated or cannot be performed (for evaluation of the extracranial nerve course) <ul style="list-style-type: none"> ○ If < 40 years of age or atypical features (i.e., bilateral, hearing loss, dizziness/vertigo, visual changes, sensory loss, numbness, pain > 2min, pain outside trigeminal nerve distribution, progression) (Policeni, 2017; Hughes, 2016; ACR CN, 2017) <p>Added:</p> <ul style="list-style-type: none"> ● Suspected infection: Abscess ● Face mass: Known or highly suspected head and neck cancer on examination ● Facial trauma: Severe facial trauma <p>Deleted:</p> <ul style="list-style-type: none"> ● Symptoms persist after four (4) consecutive weeks of medication, e.g., antibiotics, steroids or antihistamines ● Clinical Suspicion of osteomyelitis: Direct visualization of lesion over bone
--	---

	<p>Deleted:</p> <ul style="list-style-type: none"> • Face Mass <ul style="list-style-type: none"> ○ Unless increased risk for malignancy based on <ul style="list-style-type: none"> ▪ Any of these: ▪ Fixation to adjacent tissues ▪ Firm consistency ▪ Size >1.5 cm ▪ Ulceration of overlying skin ○ Clinical concern for abscess • Facial trauma: Physical findings of direct facial bone injury
<p>May 2019</p>	<ul style="list-style-type: none"> • Added: Suspected orbital trauma w/indeterminate x-ray or US • Added specifics to Face Mass: <ul style="list-style-type: none"> ○ Present on physical exam and remains non-diagnostic after x-ray or ultrasound is completed (Kuno, 2014) ○ Clinical concern for abscess ○ Failed 2 weeks of treatment for suspected infectious adenopathy (Haynes, 2015). • Removed: <ul style="list-style-type: none"> ○ Hyposmia • Immunocompromised patient

REFERENCES

1. Chiarella SE, Grammer LC. Immune deficiency in chronic rhinosinusitis: screening and treatment. *Expert Rev Clin Immunol*. Feb 2017;13(2):117-123. doi:10.1080/1744666x.2016.1216790
2. Kaplan A. Canadian guidelines for chronic rhinosinusitis: Clinical summary. *Can Fam Physician*. 2013;59(12):1275-e534.
3. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngol Head Neck Surg*. Apr 2015;152(2 Suppl):S1-s39. doi:10.1177/0194599815572097
4. Brook I, Hinthorn DR. Chronic sinusitis questions & answers. WebMD, LLC. Updated July 1, 2019. Accessed November 30, 2021. <https://emedicine.medscape.com/article/232791-questions-and-answers>
5. Frerichs N, Brateanu A. Rhinosinusitis and the role of imaging. *Cleve Clin J Med*. Jul 31 2020;87(8):485-492. doi:10.3949/ccjm.87a.19092
6. American College of Radiology. ACR Appropriateness Criteria® Cranial Neuropathy. American College of Radiology (ACR). Updated 2017. Accessed November 2, 2021. <https://acsearch.acr.org/docs/69509/Narrative/>
7. Silveira MLC, Anselmo-Lima WT, Faria FM, et al. Impact of early detection of acute invasive fungal rhinosinusitis in immunocompromised patients. *BMC Infect Dis*. Apr 5 2019;19(1):310. doi:10.1186/s12879-019-3938-y
8. Dankbaar JW, van Bommel AJ, Pameijer FA. Imaging findings of the orbital and intracranial complications of acute bacterial rhinosinusitis. *Insights Imaging*. Oct 2015;6(5):509-18. doi:10.1007/s13244-015-0424-y
9. Kastner J, Simmen D, Netuka D, Kastner J, Gudziol V. Orbital and Intracranial Complications of Acute and Chronic Rhinosinusitis. In: Chang CC, Incaudo GA, Gershwin ME, eds. *Diseases of the Sinuses: A Comprehensive Textbook of Diagnosis and Treatment*. Springer New York; 2014:495-515.
10. Seidman MD, Gurgel RK, Lin SY, et al. Clinical Practice Guideline: Allergic Rhinitis. *Otolaryngol Head Neck Surg*. 2015/02/01 2015;152(1_suppl):S1-S43. doi:10.1177/0194599814561600
11. Vashishta R, Soler ZM, Nguyen SA, Schlosser RJ. A systematic review and meta-analysis of asthma outcomes following endoscopic sinus surgery for chronic rhinosinusitis. *Int Forum Allergy Rhinol*. Oct 2013;3(10):788-94. doi:10.1002/alr.21182
12. American College of Radiology. ACR Appropriateness Criteria® Sinusitis--Child. American College of Radiology. Updated 2018. Accessed November 30, 2021. <https://acsearch.acr.org/docs/69442/Narrative/>
13. Wald ER, Applegate KE, Bordley C, et al. Clinical practice guideline for the diagnosis and management of acute bacterial sinusitis in children aged 1 to 18 years. *Pediatrics*. Jul 2013;132(1):e262-80. doi:10.1542/peds.2013-1071
14. Poorey VK, Gupta N. Endoscopic and computed tomographic evaluation of influence of nasal septal deviation on lateral wall of nose and its relation to sinus diseases. *Indian J Otolaryngol Head Neck Surg*. Sep 2014;66(3):330-5. doi:10.1007/s12070-014-0726-2

15. Sedaghat AR, Kieff DA, Bergmark RW, Cunnane ME, Busaba NY. Radiographic evaluation of nasal septal deviation from computed tomography correlates poorly with physical exam findings. *Int Forum Allergy Rhinol*. Mar 2015;5(3):258-62. doi:10.1002/alr.21445
16. Kirsch CFE, Bykowski J, Aulino JM, et al. ACR Appropriateness Criteria(®) Sinonasal Disease. *J Am Coll Radiol*. Nov 2017;14(11s):S550-s559. doi:10.1016/j.jacr.2017.08.041
17. American College of Radiology. ACR Appropriateness Criteria® Sinonasal Disease. American College of Radiology (ACR). Updated 2021. Accessed March 17, 2022. <https://acsearch.acr.org/docs/69502/Narrative/>
18. Sahay S, Gera K, Bhargava SK, Shah A. Occurrence and impact of sinusitis in patients with asthma and/or allergic rhinitis. *J Asthma*. Aug 2016;53(6):635-43. doi:10.3109/02770903.2015.1091005
19. Ragab S, Scadding GK, Lund VJ, Saleh H. Treatment of chronic rhinosinusitis and its effects on asthma. *Eur Respir J*. Jul 2006;28(1):68-74. doi:10.1183/09031936.06.00043305
20. Geyer M, Nilssen E. Evidence-based management of a patient with anosmia. *Clin Otolaryngol*. Oct 2008;33(5):466-9. doi:10.1111/j.1749-4486.2008.01819.x
21. Allis TJ, Leopold DA. Smell and taste disorders. *Facial Plast Surg Clin North Am*. Feb 2012;20(1):93-111. doi:10.1016/j.fsc.2011.10.011
22. Pincus DJ, Armstrong MB, Thaller SR. Osteomyelitis of the craniofacial skeleton. *Semin Plast Surg*. 2009;23(2):73-79. doi:10.1055/s-0029-1214159
23. Koeller KK. Radiologic Features of Sinonasal Tumors. *Head Neck Pathol*. Mar 2016;10(1):1-12. doi:10.1007/s12105-016-0686-9
24. Haynes J, Arnold KR, Aguirre-Oskins C, Chandra S. Evaluation of neck masses in adults. *Am Fam Physician*. May 15 2015;91(10):698-706.
25. Echo A, Troy JS, Hollier LH, Jr. Frontal sinus fractures. *Semin Plast Surg*. 2010;24(4):375-382. doi:10.1055/s-0030-1269766
26. American College of Radiology. ACR Appropriateness Criteria® Head Trauma. American College of Radiology (ACR). Updated 2020. Accessed November 2, 2021. <https://acsearch.acr.org/docs/69481/Narrative/>
27. American College of Radiology. ACR Appropriateness Criteria® Major Blunt Trauma. American College of Radiology. Updated 2019. Accessed November 30, 2021. <https://acsearch.acr.org/docs/3102405/Narrative/>
28. Oh JW, Kim SH, Whang K. Traumatic Cerebrospinal Fluid Leak: Diagnosis and Management. *Korean J Neurotrauma*. Oct 2017;13(2):63-67. doi:10.13004/kjnt.2017.13.2.63
29. Raju N, Ishwar P, Banerjee R. Role of multislice computed tomography and three-dimensional rendering in the evaluation of maxillofacial injuries. Original Article. *Journal of Oral and Maxillofacial Radiology*. September 1, 2017 2017;5(3):67-73. doi:10.4103/jomr.jomr_25_17
30. Vemuri NV, Karanam LSP, Manchikanti V, Dandamudi S, Puvvada SK, Vemuri VK. Imaging review of cerebrospinal fluid leaks. *Indian J Radiol Imaging*. Oct-Dec 2017;27(4):441-446. doi:10.4103/ijri.IJRI_380_16
31. Abdel Razek AAK, Mukherji S. Imaging of sialadenitis. *Neuroradiol J*. Jun 2017;30(3):205-215. doi:10.1177/1971400916682752
32. Gadodia A, Bhalla AS, Sharma R, Thakar A, Parshad R. Bilateral parotid swelling: a radiological review. *Dentomaxillofac Radiol*. 2011;40(7):403-414. doi:10.1259/dmfr/17889378

33. Kalia V, Kalra G, Kaur S, Kapoor R. CT Scan as an Essential Tool in Diagnosis of Non-radiopaque Sialoliths. *J Maxillofac Oral Surg*. Mar 2015;14(Suppl 1):240-4. doi:10.1007/s12663-012-0461-8
34. Terraz S, Poletti PA, Dulguerov P, et al. How reliable is sonography in the assessment of sialolithiasis? *AJR Am J Roentgenol*. Jul 2013;201(1):W104-9. doi:10.2214/ajr.12.9383
35. Pakalniskis MG, Berg AD, Policeni BA, et al. The Many Faces of Granulomatosis With Polyangiitis: A Review of the Head and Neck Imaging Manifestations. *AJR Am J Roentgenol*. Dec 2015;205(6):W619-29. doi:10.2214/ajr.14.13864
36. Popovic KS, Kocar M. Imaging findings in bisphosphonate-induced osteonecrosis of the jaws. *Radiol Oncol*. Dec 2010;44(4):215-9. doi:10.2478/v10019-010-0032-x
37. Andreu-Arasa VC, Chapman MN, Kuno H, Fujita A, Sakai O. Craniofacial Manifestations of Systemic Disorders: CT and MR Imaging Findings and Imaging Approach. *RadioGraphics*. 2018/05/01 2018;38(3):890-911. doi:10.1148/rg.2018170145
38. Borges A, Casselman J. Imaging the trigeminal nerve. *Eur J Radiol*. May 2010;74(2):323-40. doi:10.1016/j.ejrad.2010.02.006
39. Lohrmann C, Uhl M, Warnatz K, Kotter E, Ghanem N, Langer M. Sinonasal computed tomography in patients with Wegener's granulomatosis. *J Comput Assist Tomogr*. Jan-Feb 2006;30(1):122-5. doi:10.1097/01.rct.0000191134.67674.c6
40. Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol*. Aug 2020;277(8):2251-2261. doi:10.1007/s00405-020-05965-1
41. Saniasiaya J, Islam MA, Abdullah B. Prevalence of Olfactory Dysfunction in Coronavirus Disease 2019 (COVID-19): A Meta-analysis of 27,492 Patients. *Laryngoscope*. Apr 2021;131(4):865-878. doi:10.1002/lary.29286
42. Phal PM, Myall RW, Assael LA, Weissman JL. Imaging findings of bisphosphonate-associated osteonecrosis of the jaws. *AJNR Am J Neuroradiol*. Jun-Jul 2007;28(6):1139-45. doi:10.3174/ajnr.A0518
43. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. Jan 2018;38(1):1-211. doi:10.1177/0333102417738202

ADDITIONAL RESOURCES

1. Abuabara A. Cerebrospinal fluid rhinorrhoea: diagnosis and management. *Med Oral Patol Oral Cir Bucal*. Sep 1 2007;12(5):E397-400.
2. Borges A, Casselman J. Imaging the trigeminal nerve. *Eur J Radiol*. May 2010;74(2):323-40. doi:10.1016/j.ejrad.2010.02.006
3. De Lucas EZ, Martínez JF-L, Ribate DR, et al. Analysis of the findings on paranasal sinuses and chest computed tomography in patients with severe asthma. *Eur Respir J*. 2018;52:PA677. doi:10.1183/13993003.congress-2018.PA677

4. Decker JR, Meen EK, Kern RC, Chandra RK. Cost effectiveness of magnetic resonance imaging in the workup of the dysosmia patient. *Int Forum Allergy Rhinol*. Jan 2013;3(1):56-61. doi:10.1002/alr.21066
5. Harreld JH, Kaufman RA, Kang G, et al. Utility of Pre-Hematopoietic Cell Transplantation Sinus CT Screening in Children and Adolescents. *AJNR Am J Neuroradiol*. May 2020;41(5):911-916. doi:10.3174/ajnr.A6509
6. Jang A-S. The role of rhinosinusitis in severe asthma. *Korean J Intern Med*. 2013;28(6):646-651. doi:10.3904/kjim.2013.28.6.646
7. Mantur M, Łukaszewicz-Zajac M, Mroczko B, et al. Cerebrospinal fluid leakage--reliable diagnostic methods. *Clin Chim Acta*. May 12 2011;412(11-12):837-40. doi:10.1016/j.cca.2011.02.017
8. Moeller CW, Martin J, Welch KC. Sinonasal evaluation preceding hematopoietic transplantation. *Otolaryngol Head Neck Surg*. May 2011;144(5):796-801. doi:10.1177/0194599810395089
9. Setzen G, Ferguson BJ, Han JK, et al. Clinical consensus statement: appropriate use of computed tomography for paranasal sinus disease. *Otolaryngol Head Neck Surg*. Nov 2012;147(5):808-16. doi:10.1177/0194599812463848
10. Varshney H, Varshney J, Biswas S, Ghosh SK. Importance of CT Scan of Paranasal Sinuses in the Evaluation of the Anatomical Findings in Patients Suffering from Sinonasal Polyposis. *Indian J Otolaryngol Head Neck Surg*. Jun 2016;68(2):167-72. doi:10.1007/s12070-015-0827-6

Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: National Imaging Associates, Inc. (NIA) authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Evolent Health LLC subsidiaries including, but not limited to, National Imaging Associates (“NIA”). The policies constitute only the reimbursement and coverage guidelines of NIA. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. NIA reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.