



Evoleant*	
Clinical guidelines PLAIN FILM X-RAYS	Original Date: April 2016
Physical Medicine – Clinical Decision Making	Last Revised Date: August 2022
Guideline Number: Evoleant_CG_610	Implementation Date: July 2023

Policy Statement

The use of plain films is medically necessary when clinical findings dictate their utilization. Films are not indicated to identify unsuspected contraindications to chiropractic manipulation, view postural changes and biomechanics, or identify subluxations. Insufficient scientific evidence exists to support the use of routine plain film radiographs as a means for improved clinical outcomes in spinal disorders. There is insufficient clinical research to support improved clinical outcomes when radiographs are a part of a routine component of the initial evaluation or ongoing treatment. This organization has adopted the Diagnostic Imaging Practice Guidelines for Musculoskeletal Complaints in Adults.¹⁻⁴ These guidelines represent the official position of the Council on Chiropractic Guidelines and Practice Parameters in matters related to the use of diagnostic imaging in the chiropractic profession.

The use of full spine radiographs, except for the clinical investigation and diagnosis of scoliosis, is not supported by clinical research.

Initial Clinical Reviewers (ICRs) and Physician Clinical Reviewers (PCRs) must be able to apply criteria based on individual needs and based on an assessment of the local delivery system.

Purpose

This policy will be used to support the medical necessity of plain film radiographs by chiropractic providers within the first 30 days of care.

Scope

This policy will apply to all participating network chiropractic practitioners.

Definition

Plain films:

Spinal or extremity radiographs used as a diagnostic tool by chiropractors.

* Evoleant refers to Evoleant Health LLC and Evoleant Specialty Services, Inc.

Guidelines

- An appropriate history and examination are required to determine if plain films are clinically indicated.
- Utilization of radiographs by chiropractors will not be reimbursed unless sufficient medical record documentation is submitted with claims to support the medical necessity of the film. The clinical record must clearly document the rationale for the x-rays, any suspected pathology, or what condition the chiropractor hopes to rule out. The use of plain films to rule out an unsuspected pathology is not clinically indicated.
- Routine use of radiographs as part of the initial evaluation or part of an ongoing treatment plan will not be reimbursed.
- The use of full spine radiographs for any diagnosis other than scoliosis is not considered medically necessary and will not be reimbursed.
- Contraindications to plain film x-rays include:
 - Infants (0 - 36 months)
 - Pregnancy or possible pregnancy
 - Obesity, if size precludes good radiographic resolution
 - Patient has positioning difficulty due to mental status or physical restrictions, which precludes good radiographic resolution
 - Children 3 to 18 years of age, except for investigation of suspected acute fracture, dislocation, infection, scoliosis, developmental defects, or a suspected pathology.

Requirements

- The clinical record must contain a written x-ray report within 5 business days from the date of service.
- The clinic must have all of the following:
 - A documented Quality Control Program inclusive of both imaging equipment and film processors
 - A documented Radiation Safety and As Low As Reasonably Achievable (ALARA) Program
 - Documented emergency policies, procedures, and equipment on site (i.e., automated external defibrillator (AED))
 - Documentation of current Basic Life Support (BLS) certification
 - Records of formal preventative maintenance program per original equipment specifications
 - A current (within 3 years) letter of state inspection, calibration report, or physicist's report

- At a minimum, an automatic processor must be used to develop all analog plain films.

Clinical Examples of Medically Necessary X-rays

- Investigation of suspected acute fracture
- Follow-up radiographs to monitor a healing fracture
- Investigation of suspected bony dislocation
- Evaluation of prior surgical site where manual based treatment may be applied (where no previous films are available for review)
- Suspected (patient history, pain characteristics and/or physical examination) malignancy, infection, systemic disease, or inflammatory spondyloarthropathy
- Precise quantification of clinically suspected active child or juvenile scoliosis
- Persistent (same or worse pain) after first month of treatment
- Significant history of drug or alcohol abuse, such as IV drugs, chronic alcoholism, or chronic use of steroids
- Adult with thoracolumbar, lumbar, or thoracic spine blunt trauma or acute injuries (falls, motor vehicle accidents [MVA], motorcycle, pedestrian, cyclists, etc.)
- Adult with complicated (i.e., “red flag”) low back pain (LBP), thoracic pain, or neck pain **and** indicators of contraindication to spinal manipulative therapy (SMT) (relative/absolute)
- Suspected inflammatory spondyloarthropathies⁵, neoplasia, or infection
- Adult: in the absence of expected treatment response or worsening after 4 weeks of conservative treatment
- Adult with acute neck injury and positive CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)
- Suspected lumbar degenerative spinal stenosis or spondylolisthesis if individual is greater than 50 years of age and/or has progressive neurological deficit – AP (or PA) and lateral lumbar views⁴
- Adult with recent unimaged blunt trauma to pelvis and unable to bear weight¹ – AP pelvis and lateral hip “frog leg” views⁴
- Acute neck pain with recent unimaged dangerous trauma; paresthesia in extremities; age greater than 65 years; or non-traumatic neck pain with radicular symptoms¹ – APOM, AP lower cervical and lateral neutral views⁴
- Adult with painful or progressive scoliosis¹ – Erect standing full spine (14x36) PA and lateral views in the absence of recent films.⁴
- Plain film x-rays may be appropriate when red flags suggest further screening for cancer, infection, or fracture. They may also be sufficient for the initial evaluation of individuals with the following red flags: age >70 years, a history of recent significant trauma, or risk of osteoporosis. Plain film x-rays may be appropriate but are usually not sufficient for clinical decision making without advanced imaging in the presence of other red flags. Radiographs are unreliable for assessment of bone mass changes before at least a 30%-50% loss. In healthy peri- and early postmenopausal women (age 45-64), consider using the Osteoporosis Self-Assessment Tool (OST score). The OST score considers only 2

variables: $(\text{weight in kg} - \text{age})/5$. The cut-off for a positive test is <2 , indicating this woman should be referred for DXA.⁴

- Current x-ray recommendations/guidelines for spinal and extremity disorders emphasize a focused history and physical examination, reassurance, initial pain management medications if necessary (acetaminophen or nonsteroidal anti-inflammatory drugs), and consideration of nonpharmacologic therapies (e.g., manipulation, exercise, etc.) without routine imaging in individuals with nonspecific neck and/or low back pain. Imaging is considered for those without improvement after 6 weeks and for those with clinical indicators of serious pathologies (red flags).

Plain film x-rays of the extremities may be indicated in the following circumstances:

- Significant history of recent trauma sufficient to cause fracture
- Significant history of repetitive stress to cause stress fracture
- Suspected stress (insufficiency) fracture¹³
- History or clinical findings of malignancy
- Previous surgery or fracture
- Suspicion of or confirmed inflammatory arthritis
- Evaluation of gross deformities
- Bruising, swelling, redness heat, indicating infection
- Lymphadenopathy
- Evaluation of developmental hip dysplasia in the pediatric population
- Evaluation of Leg-Calve-Perthes disease
- Evaluation of slipped capital femoral epiphysis in the pediatric population
- Evaluation of chronic hip pain, initial imaging

Plain film radiographs may be appropriate but are usually not sufficient for clinical decision making without advanced imaging (MR and/or CT) in the presence of other red flags including:

- Age < 20 years or > 50 years
- Failure to improve with care, no prior films
- Personal history of intravenous drug abuse
- History of malignancy
- Immune suppression
- Night pain (including when unrelated to movement)
- Pain at multiple sites
- Pain at rest
- Fever
- Structural deformity
- Systemic unwellness
- Unexplained weight loss

Spinal radiographs also have a role in evaluation of scoliosis and in postoperative evaluation of instrumentation and fusion. For the evaluation of scoliosis in children, radiographic decision-making and examinations should be performed in accordance with guidance published by the

American College of Radiology (ACR) and the Society for Pediatric Radiology (SPR). Radiographic examination is indicated for pediatric patients at high risk for cervical spine instability – especially those with Down syndrome.¹⁷

Initial Plain Film X-rays Are Not Indicated in the Following Cases

- Adult with acute uncomplicated pain (< 4 weeks duration) in any of the following
 - LBP¹ (uncomplicated definition: nontraumatic pain without neurologic deficits or indicators of potentially serious pathologies)⁴
 - Thoracic spine pain^{1,4}
 - Uncomplicated neck pain^{1,4}
- Adult with nontraumatic acute LBP (<4 weeks duration) AND sciatica and no red flags^{1,4}
- Adult with uncomplicated subacute pain (4-12 weeks duration) in any of the following
 - LBP and no previous treatment trial^{1,4}
 - Thoracic spine pain and no previous treatment trial^{1,4}
 - Subacute neck pain with or without arm pain^{1,4}
- Adult with persistent pain (>12 weeks) in any of the following
 - LBP and no previous treatment trial^{1,4}
 - Thoracic spine pain and no previous treatment trial^{1,4}
 - Persistent neck pain with or without arm pain^{1,4}
- Sciatica, unless individual is age >50 or has progressive neurological deficits⁴
- Suspected lumbar disc herniation
- Suspected degenerative spondylolithesis/lateral stenosis, unless individual is age >50 or has progressive neurological deficits
- Suspected lumbar degenerative spinal stenosis, unless individual is age >50 or has progressive neurological deficits
- Adult with nonpainful and nonprogressive scoliosis
- Adult with acute neck injury and negative CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)
- In headache complaints, vital signs (to R/O severe hypertension or fever) and testing of the cranial nerves (to R/O vascular events, space occupying lesions, etc.) should be an integral part of initial examination. Significant positive findings mandate further evaluation.⁴ Without red flags or significant findings, no initial films are indicated.
- Coccyx trauma and coccydynia
- The routine use of spinal radiographs for structural and biomechanical analysis has not been substantiated to improve patient outcomes.^{18 19} The clinical evidence is insufficient to support an association between sagittal (lordosis, kyphosis) spinal curves and health outcomes including spine-related pain.²⁰ The utility of plain film radiography for the detection of spinal ‘subluxations’, or to guide the specifics of spinal manipulative therapy, is controversial.¹⁸ “The validity of the various systems of roentgenometric analysis has not been proven and their underlying premise of bilateral symmetry within the body does not take into account natural structural anomalies”.¹⁸ Adding to this

controversy is the fact that nonspecific spinal abnormalities are common in asymptomatic patients.

- Chou et al state that “[s]trong evidence shows that routine back imaging does not improve patient outcomes, exposes patients to unnecessary harms, and increases costs”.²¹ Further, Andersen notes that “[a]vailable evidence indicates that immediate, routine lumbar spine imaging in patients with LBP and without features indicating a serious underlying condition, did not improve outcomes compared with usual clinical care without immediate imaging. Clinical care without immediate imaging seems to result in no increased odds of failure in identifying serious underlying conditions in patients without risk factors for these conditions. In addition to lacking clinical benefit, routine lumbar imaging is associated with radiation exposure (radiography and CT) and increased direct expenses for patients and may lead to unnecessary procedures. This evidence confirms that clinicians should refrain from routine, immediate lumbar imaging in primary care patients with nonspecific, acute or subacute LBP and no indications of underlying serious conditions.”²² Among the recommendations given for Chiropractic Management of Patients with Chronic Musculoskeletal Pain, Hawk et al, recommends avoiding the use of routine imaging, “because chronic MSK pain is often multifactorial and may not originate from a local source, imaging evidence is rarely capable of definitively identifying a pain source. However, imaging may be necessary if red flags are present and should be evaluated on a case-by-case basis after a thorough history and examination are performed.”²³

POLICY HISTORY

Date	Summary
August 2022	<ul style="list-style-type: none"> • Rearranged criteria under “Initial Plain Film X-rays Are Not Indicated in the Following Cases”- content was not changed • Added under plain film x-rays of the extremities <ul style="list-style-type: none"> ○ Evaluation of chronic hip pain – initial imaging ○ Suspected stress (insufficiency) fracture
November 2021	<ul style="list-style-type: none"> • Added “General Information” statement • Clarified intent of the following indication by changing the format of ‘and’: <ul style="list-style-type: none"> ○ “Adults with complicated (i.e., “red flag”) LBP, thoracic pain, or neck pain and indicators of contraindication to SMT (relative/absolute)”
October 2020	No content changes
January 2020	No edits made to guideline in response to the review of the evidence base
July 2019	Format updated for improved readability

REFERENCES

1. Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults--an evidence-based approach: introduction. *J Manipulative Physiol Ther.* Nov-Dec 2007;30(9):617-83. doi:10.1016/j.jmpt.2007.10.003
2. Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging guideline for musculoskeletal complaints in adults-an evidence-based approach-part 2: upper extremity disorders. *J Manipulative Physiol Ther.* Jan 2008;31(1):2-32. doi:10.1016/j.jmpt.2007.11.002
3. Bussièrès AE, Taylor JA, Peterson C. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults--an evidence-based approach. Part 1. Lower extremity disorders. *J Manipulative Physiol Ther.* Nov-Dec 2007;30(9):684-717. doi:10.1016/j.jmpt.2007.10.004
4. Bussièrès AE, Taylor JA, Peterson C. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults-an evidence-based approach-part 3: spinal disorders. *J Manipulative Physiol Ther.* Jan 2008;31(1):33-88. doi:10.1016/j.jmpt.2007.11.003
5. Czuczman GJ, Mandell JC, Wessell DE, et al. ACR Appropriateness Criteria® Inflammatory Back Pain: Known or Suspected Axial Spondyloarthritis: 2021 Update. *J Am Coll Radiol.* Nov 2021;18(11s):S340-s360. doi:10.1016/j.jacr.2021.08.003
6. Stiell IG, Wells GA, Vandemheen KL, et al. The Canadian C-Spine Rule for Radiography in Alert and Stable Trauma Patients. *JAMA.* 2001;286(15):1841-1848. doi:10.1001/jama.286.15.1841
7. Dagenais S, Tricco A, Haldeman S. Assessment of Low Back Pain. In: Dagenais S, Haldeman S, eds. *Evidence-Based Management of Low Back Pain.* Mosby (Elsevier) 2012:21-31.
8. Koes BW, van Tulder M, Lin C-WC, Macedo LG, McAuley J, Maher C. An updated overview of clinical guidelines for the management of non-specific low back pain in primary care. *Eur Spine J.* 2010;19(12):2075-2094. doi:10.1007/s00586-010-1502-y
9. Pillastrini P, Gardenghi I, Bonetti F, et al. An updated overview of clinical guidelines for chronic low back pain management in primary care. *Joint Bone Spine.* Mar 2012;79(2):176-85. doi:10.1016/j.jbspin.2011.03.019
10. Bach SM, Holten KB. Guideline update: what's the best approach to acute low back pain? *J Fam Pract.* Dec 2009;58(12):E1.
11. Chou R, Qaseem A, Owens DK, Shekelle P. Diagnostic imaging for low back pain: advice for high-value health care from the American College of Physicians. *Ann Intern Med.* Feb 1 2011;154(3):181-9. doi:10.7326/0003-4819-154-3-201102010-00008
12. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* Oct 2 2007;147(7):478-91. doi:10.7326/0003-4819-147-7-200710020-00006
13. Expert Panel on Musculoskeletal I, Bencardino JT, Stone TJ, et al. ACR Appropriateness Criteria((R)) Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *J Am Coll Radiol.* May 2017;14(5S):S293-S306. doi:10.1016/j.jacr.2017.02.035
14. Davis PC, Wippold FJ, 2nd, Brunberg JA, et al. ACR Appropriateness Criteria on low back pain. *J Am Coll Radiol.* Jun 2009;6(6):401-7. doi:10.1016/j.jacr.2009.02.008
15. Hutchins TA, Peckham M, Shah LM, et al. ACR Appropriateness Criteria® Low Back Pain. American College of Radiology. Updated 2021. Accessed August 15, 2022.

16. American College of Radiology, Society for Pediatric Radiology, Society of Skeletal Radiology. ACR-SPR-SSR practice parameter for the performance of radiography for scoliosis in children. American College of Radiology (ACR). Updated 2019. Accessed August 15, 2022.
17. American College of Radiology, American Society of Spine Radiology, Society for Pediatric Radiology, Society of Skeletal Radiology. ACR-ASSR-SPR-SSR practice parameter for the performance of spine radiography. American College of Radiology (ACR). Updated 2022. Accessed August 30, 2022.
18. Petersen C, Hsu W. Indications for and use of x-rays. *Principles and Practice of Chiropractic*. 3rd ed. McGraw-Hill Education; 2005:661-681.
19. Corso M, Cancelliere C, Mior S, Kumar V, Smith A, Côté P. The clinical utility of routine spinal radiographs by chiropractors: a rapid review of the literature. *Chiropr Man Therap*. Jul 9 2020;28(1):33. doi:10.1186/s12998-020-00323-8
20. Christensen ST, Hartvigsen J. Spinal curves and health: a systematic critical review of the epidemiological literature dealing with associations between sagittal spinal curves and health. *J Manipulative Physiol Ther*. Nov-Dec 2008;31(9):690-714. doi:10.1016/j.jmpt.2008.10.004
21. Chou R, Deyo RA, Jarvik JG. Appropriate use of lumbar imaging for evaluation of low back pain. *Radiol Clin North Am*. Jul 2012;50(4):569-85. doi:10.1016/j.rcl.2012.04.005
22. Andersen JC. Is immediate imaging important in managing low back pain? *J Athl Train*. Jan-Feb 2011;46(1):99-102. doi:10.4085/1062-6050-46.1.99
23. Hawk C, Whalen W, Farabaugh RJ, et al. Best Practices for Chiropractic Management of Patients with Chronic Musculoskeletal Pain: A Clinical Practice Guideline. *J Altern Complement Med*. Oct 2020;26(10):884-901. doi:10.1089/acm.2020.0181

ADDITIONAL RESOURCES

1. American Chiropractic Association. Avoid routine spinal imaging in the absence of clear clinical indicators for patients with acute low back pain of less than six (6) weeks duration. ABIM Foundation. Updated July 12, 2021. Accessed August 15, 2022. <https://www.choosingwisely.org/clinician-lists/aca-spinal-imaging-for-acute-low-back-pain/>
2. Ammendolia C, Hogg-Johnson S, Pennick V, Glazier R, Bombardier C. Implementing evidence-based guidelines for radiography in acute low back pain: a pilot study in a chiropractic community. *J Manipulative Physiol Ther*. Mar-Apr 2004;27(3):170-9. doi:10.1016/j.jmpt.2003.12.021
3. Acute low back problems in adults: assessment and treatment. Agency for Health Care Policy and Research. *Clin Pract Guidel Quick Ref Guide Clin*. Dec 1994;(14):iii-iv, 1-25.
4. Borenstein DG. A clinician's approach to acute low back pain. *Am J Med*. Jan 27 1997;102(1a):16s-22s. doi:10.1016/s0002-9343(97)00412-9
5. Dagenais S, Tricco AC, Haldeman S. Synthesis of recommendations for the assessment and management of low back pain from recent clinical practice guidelines. *Spine J*. Jun 2010;10(6):514-29. doi:10.1016/j.spinee.2010.03.032
6. Henschke N, Maher CG, Ostelo RW, de Vet HC, Macaskill P, Irwig L. Red flags to screen for malignancy in patients with low-back pain. *Cochrane Database Syst Rev*. Feb 28 2013;(2):Cd008686. doi:10.1002/14651858.CD008686.pub2

7. Institute for Clinical Systems Improvement. Health care guideline: Adult acute and subacute low back pain. Institute for Clinical Systems Improvement (ICSI). Updated March 2018. Accessed August 15, 2022. <https://www.icsi.org/wp-content/uploads/2019/08/March-2018-LBP-Interactive2.pdf>
8. Jarvik JG. Imaging of adults with low back pain in the primary care setting. *Neuroimaging Clin N Am*. May 2003;13(2):293-305. doi:10.1016/s1052-5149(03)00022-4
9. Kendrick D, Fielding K, Bentley E, Kerslake R, Miller P, Pringle M. Radiography of the lumbar spine in primary care patients with low back pain: randomised controlled trial. *BMJ (Clinical research ed)*. 2001;322(7283):400-405. doi:10.1136/bmj.322.7283.400
10. Kendrick D, Fielding K, Bentley E, Miller P, Kerslake R, Pringle M. The role of radiography in primary care patients with low back pain of at least 6 weeks duration: a randomised (unblinded) controlled trial. *Health Technol Assess*. 2001;5(30):1-69. doi:10.3310/hta5300
11. Magora A. Relation between the low back pain syndrome and x-ray findings. Degenerative osteoarthritis. *Scand J Rehabil Med*. 1976;8:115-125.
12. Mootz R, Hoffman L, Hansen D. Optimizing clinical use of radiography and minimizing radiation exposure in chiropractic practice. *Topics in Clinical Chiropractic*. 1997;4(1):34-44.
13. (NASS) NASS. *Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis & Treatment of Low Back Pain*. North American Spine Society. Updated 2020. Accessed November 4, 2022. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.pdf>
14. New Zealand Guidelines Group, New Zealand Society of Physiotherapists, Royal New Zealand College of General Practitioners, New Zealand Register of Osteopaths. New Zealand Acute Low Back Pain Guide. Updated October 2004. Accessed August 15, 2022. <https://www.healthnavigator.org.nz/media/1006/nz-acute-low-back-pain-guide-acc.pdf>
15. Patel ND, Broderick DF, Burns J, et al. ACR Appropriateness Criteria Low Back Pain. *J Am Coll Radiol*. Sep 2016;13(9):1069-78. doi:10.1016/j.jacr.2016.06.008
16. Tran B, Saxe JM, Ekeh AP. Are flexion extension films necessary for cervical spine clearance in patients with neck pain after negative cervical CT scan? *J Surg Res*. Sep 2013;184(1):411-3. doi:10.1016/j.jss.2013.05.100
17. van den Hoogen HM, Koes BW, van Eijk JT, Bouter LM. On the accuracy of history, physical examination, and erythrocyte sedimentation rate in diagnosing low back pain in general practice. A criteria-based review of the literature. *Spine (Phila Pa 1976)*. Feb 1 1995;20(3):318-27. doi:10.1097/00007632-199502000-00010
18. van Tulder MW, Tuut M, Pennick V, Bombardier C, Assendelft WJ. Quality of primary care guidelines for acute low back pain. *Spine (Phila Pa 1976)*. Sep 1 2004;29(17):E357-62. doi:10.1097/01.brs.0000137056.64166.51
19. Williams CM, Henschke N, Maher CG, et al. Red flags to screen for vertebral fracture in patients presenting with low-back pain. *Cochrane Database Syst Rev*. Jan 31 2013;(1):Cd008643. doi:10.1002/14651858.CD008643.pub2
20. Wipf JE, Deyo RA. Low back pain. *Med Clin North Am*. Mar 1995;79(2):231-46. doi:10.1016/s0025-7125(16)30065-7
21. Yelland M. Diagnostic imaging for back pain. *Aust Fam Physician*. Jun 2004;33(6):415-9.

22. Young KJ. Historical influence on the practice of chiropractic radiology: Part I - a survey of Diplomates of the American Chiropractic College of Radiology. *Chiropr Man Therap.* 2017;25:14-14. doi:10.1186/s12998-017-0146-y

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Reviewed / Approved by Clinical Guideline Committee

Disclaimer: Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. **Members should contact their Plan customer service representative for specific coverage information.**